

**Fortegnelse, kontroldokument
og brugervejledning kategori A – MFAG A2**

**Inventory, Control Document
and User Instructions Category A –
MFAG A2**



Fortegnelse, kontroldokument og brugervejledning over lægemiddelstoffer og medicinsk udstyr kategori A- MFAG A2

Jf. Meddelelser fra Søfartsstyrelsen A, Kapitel IX B

Inventory, Control Document, User Instructions of Active Pharmaceutical Ingredients and Medical Equipment Category A – MFAG A2

Cf. Danish Maritime Authority Notice A, chapter IX B

July 2024

ISBN: 978-87-7454-212-4

Identifikation af skib / Identification of ship

Skibsnavn / <i>Ship Name:</i>	
-------------------------------	--

Kaldesignal / <i>Call Signal:</i>	
-----------------------------------	--

IMO nummer / <i>IMO number:</i>	
---------------------------------	--

Hjemmehavn / <i>Port of Reg.:</i>	
-----------------------------------	--

Medicin certifikat / *Medicine Certificate*

Dette dokument anses for at udgøre skibets medicin certifikat.

Det attesteres herved, at medicin og medicinsk udstyr er eftersat og fundet i overensstemmelse med Den Danske Søfartsstyrelses krav jf. Meddelelser A Kapitel IX B.

This document is considered to represent the ships certificate of medicine.

It certifies that medicine and medical equipment has been inspected and found in compliance with the Danish Maritime Authority's requirement cf. Notes A chapter IX B.

Skibsnavn / *Ship Name*:

Kaldesignal / *Call Signal*:

IMO nummer / *IMO Number*:

Hjemmehavn / *Port of Reg.*:

Dato / <i>Date</i>	Rank / <i>Rank</i>	Navn / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / <i>Master</i>		
Næste eftersyn / <i>Next Inspection</i>			
Dato / <i>Date</i>	Rank / <i>Rank</i>	Navn / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / <i>Master</i>		

Print denne side / *Print this page*

Certifikat for genoplivningsudstyr (A.1)

Certificate resuscitation equipment (A.1)

Det attesteres herved at det **mobile genoplivningsudstyr (A.1)** er efterset og fundet i overensstemmelse med Den Danske Søfartsstyrelses krav jf. Meddelelser A Kapitel IX B.

It certifies that the **mobile resuscitation equipment (A.1)** has been inspected and found in compliance with the Danish Maritime Authority's requirement cf. Notes A chapter IX B.

Skibsnavn / *Ship Name*:

Kaldesignal / *Call Signal*:

IMO nummer / *IMO Number*:

Hjemmehavn / *Port of Reg.*:

Dato / <i>Date</i>	Rank / <i>Rank</i>	Navn / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / <i>Master</i>		
Næste eftersyn / <i>Next Inspection</i>			
Dato / <i>Date</i>	Rank / <i>Rank</i>	Navn / <i>Name</i>	Underskrift / <i>Signature</i>
	Skibsfører / <i>Master</i>		

Print denne side / *Print this page*

Indhold / Contents

Brugervejledning:	5
<i>User instruction:</i>	5
Anvendte forkortelser	9
<i>Used abbreviations</i>	9
Brugervejledning Lægemiddelstoffer	10
<i>User instruction Activ Phamaceutical</i>	10
Fortegnelse og kontroldokument	65
<i>Inventory and Control Document</i>	65
Besætningens medbragte medicin	72
Fortegnelse og Kontrolldokument	74
<i>Inventory and Control Document</i>	74
Labels	83
INDEX	88
Rekvisition lægemiddelstoffer	93
<i>Requisition Active Phamaceutecal</i>	93
Ændringer	106

De Danske Medicinkister

Medicinkisten angiver medicin/medicinsk udstyr, som Radio Medical Danmark og Søfartsstyrelsen har foretrukket indenfor de rammer som EU-direktiv 92/29/EØF og den Danske lægemiddelstyrelse har givet.

Guide til:

Brugervejledning:

Første kolonne beskriver de indikationer, lægemiddelstoffet er beregnet til. De samme indikationer kan slås op i indekset bagerst i hæftet. Første kolonne beskriver også om et lægemiddelstof skal anvendes sammen med andre lægemiddelstoffer. I sådanne tilfælde er der lavet henvisninger til de relevante lægemiddelstoffer.

I samme kolonne gøres opmærksom på, om lægemiddelstoffet udgør en del af MFAG (Medical First Aid Guide), der refereres i så fald til den relevante MFAG tabel.

MFAG lægemiddelstofferne er defineret i IMDG (International Maritime Dangerous Goods) koden, supplementsdelen, seneste version og indarbejdet i denne version af den danske "Fortegnelse, kontroldokument og brugervejledning". kontroldokument og brugervejledning".

Bemærkning: Sejlads med farligt gods jævnt før IBC koden, INF koden og IGS koden er ikke indarbejdet i denne publikation. Der henvises til disse publikationer.

Anden kolonne oplyser, lægemiddelstoffets hoved- og undergruppenummer

Tredje og fjerde kolonne indeholder rækker med oplysninger om:

- **Lægemiddelstoffet:** Beskriver det aktive stof i lægemidlet. Som et eksempel kan nævnes 3.1 Paracetamol. Det forhandles over hele jorden under mange forskellige handelsnavne. Derfor kan man uanset forskellige handelsnavne, altid finde lægemidlet, hvis man kender lægemiddelstoffet.
- **Form:** Beskriver om lægemiddelstoffet skal findes som kapsel, tablet, suppositorier, injektionsvæske, infusionsvæske, creme, salve, afdelte pulver, mund spray, brusetablet, klyksma, oral opløsning eller inhalation. Form beskriver også styrken af det enkelte lægemiddelstof pr ml eller per tablet.
- **Virkning** beskriver den forventende virkning af det enkelte lægemiddel.
- **Dosering** beskriver den anbefalede dosis ved de tilstande, der er beskrevet i kolonnen indikation. Her beskrives ligeledes lægemidlets form og administrationsmåde.
Bemærkning: Radio Medical kan ordinere andre doser og administrations måder.
- **Bivirkninger** nævner de mest almindelige bivirkninger beskrevet i Dansk Lægemiddelinformation, www.medicin.dk
- **Holdbarhed** beskriver holdbarhedstiden fra produktions datoen. Holdbarhedstiden beskrevet i denne fortegnelse er fremkommet ud fra det danske handelsnavn med længst holdbarhed.
Bemærkning: Der tages forbehold for at holdbarhedstiden kan ændres.
- **Opbevaring** beskriver om der er særlige krav til opbevaring. Lægemidler skal generelt opbevares tørt og beskyttet mod lys. De fleste lægemidler tåler "almindelig temperatur" defineret som normal stuetemperatur op til 22-24° C.

The Danish Medicine Chest's

The medicine chest indicates the Active Pharmaceutical Ingredients / medical equipment that Radio Medical Denmark and Danish Maritime Authority have preferred within the framework of the EU Directive 92/29 / EEC and the Danish Medicines Agency.

Guide to:

User instruction:

First column describes the indication the Active Pharmaceutical Ingredients is intended for. The same indications can be looked up in the index at the back of the booklet. First column also describes whether an Active Pharmaceutical Ingredient should be used together with other Active Pharmaceutical Ingredients. In such cases, a reference is made to the relevant Active Pharmaceutical Ingredient.

In the same column it is noted if the Active Pharmaceutical Ingredients is a part of MFAG (Medical First Aid Guide), and reference to the relevant MFAG table is made.

MFAG Active Pharmaceutical Ingredients s are defined in the IMDG (International Maritime Dangerous Goods) code, supplement component, latest edition, and incorporated in this version of the Danish "Inventory, control document and user manual".

Note: Sailing with dangerous goods as per the IBC code, INF code and the IGC code are not included in this publication. Please refer to these publications.

Second column describes the Active Pharmaceutical Ingredient main and subgroup numbers.

Third and fourth column contains a number of information about:

- **The active pharmaceutical ingredients:** describe the Active Pharmaceutical Ingredients in the medicine. An example is 3.1 Paracetamol, which is sold all over the world under many different trade names. Regardless of different trade names it is always possible to locate the medicine, if you know the active pharmaceutical ingredients.
- **Form:** describes if the Active Pharmaceutical Ingredients is available as capsules, tablets, suppositories, injection liquid, infusion liquid, lotion, ointment, divisional powder, mouth spray, effervescent tablets, enema, oral solution or inhalation. Form also describes the strength of each active pharmaceutical ingredient per ml or per tablet.
- **Effect** describes the expected effect of each active pharmaceutical ingredients.
- **Dosage** describes the recommended dose for the condition described in the column indication. The form of the Active Pharmaceutical Ingredients as well as the usual administration is mentioned.
Note: Radio Medical may prescribe different doses and ways of administration.
- **Side effects** describes the most common side effects mentioned in the Danish Drug Information, www.medicin.dk. There is currently no English version.
- **Durability** describes the time of durability from date of production. The durability mentioned in this record is derived from the Danish trade name with the longest durability.
Note: Durability may be subject to change.
- **Storage** describes if special storage precautions are needed. The Active Pharmaceutical Ingredients should generally be kept dry and protected from light. Most medicaments tolerate "normal temperature" defined as normal room temperature up to 22-24° C.

- **Anmærkninger** beskriver, om der bør søges samråd med Radio Medical Danmark inden anvendelse af lægemiddelstoffet. Desuden beskrives, om der er specielle forholdsregler til det specifikke lægemiddelstof.

Advarsel: Lægemiddelstoffer, der har været opbevaret forkert, eller er blevet for gamle, kan være sundhedsskadelige.

Guide til:

Fortegnelse og kontrol dokument lægemiddelstoffer.

Publikationen indeholder en fortegnelse over de lægemiddelstoffer, der skal være ombord.

Rækkerne

Lægemiddelstofferne er organiseret i hovedgrupper efter deres hovedanvendelsesområde. Eksempelvis er alle smertestillende lægemiddelstoffer organiseret i gruppe 3.

Kolonnerne

Første kolonne beskriver hvert enkelt lægemiddelstoffs nummer. Eksempelvis har morfin nummeret 3.4, hvor tallet før punktum referer til hovedgruppen, og tallet efter refererer til det specifikke lægemiddelstof. Lægemiddelstofferne i skibets hospital skal så vidt muligt organiseres efter dette nummersystem

Anden kolonne beskriver ATC koden, et klassifikations system for medicin. Se mere på www.whocc.no

Tredje kolonne beskriver lægemiddelstoffet.

Bemærkning: Efter lægemiddelstoffet kan være anført et "Δ", dersom produktet er sløvende, og et "K", dersom produktet skal opbevares i køleskab.

Fjerde kolonne beskriver under "Adm.form", om lægemiddelstoffet skal findes som kapsel, tablet, suppositorier, injektionsvæske, infusionsvæske, creme, salve, afdelte pulver, mund spray, brusetablet, klyasma, oral opløsning eller inhalation.

Femte kolonne beskriver styrken pr tablet eller pr. ml.

Sjette kolonne beskriver mindste mængde, skibet skal medbringe. Skibet må gerne medbringe større mængder end det beskrevne.

Syvende kolonne MFAG A2 beskriver det supplement, skibet skal medbringe, dersom det sejler med A-fortegnelse OG har en evakuerings tid på under 24 timer.

Bemærkning: Kolonnerne mærket MFAG beskriver de minimumsmængder af supplerende lægemidler som skibet skal medbringe, dersom skibet sejler med farligt gods.

Remarks describe whether it is recommended to seek advice from Radio Medical Denmark before using the active pharmaceutical ingredients.

In addition, precautions needed for the specific Active Pharmaceutical Ingredients are described.

Warning: Active Pharmaceutical Ingredients, which have been stored improperly or are too old can be harmful to health.

Guide to:

Inventory and Control Document of Active Pharmaceutical Ingredients.

The publication contains an inventory of mandatory Active Pharmaceutical Ingredients on board.

The rows

The Active Pharmaceutical Ingredients are organized into main groups according to field of application. As an example all analgesics are organized in group 3.

The columns

First column describes each individual Active Pharmaceutical Ingredient by number. For example, morphine number 3.4, where the number before the dot refers to the main group and the number after the dot refers to the specific active pharmaceutical ingredients. The Active Pharmaceutical Ingredients in the ship's hospital must as far as possible be organized after this numbering system.

Second column describe the ATC code, a classification system on medicine. See more on www.whocc.no.

Third column describes the active pharmaceutical ingredients

Note: The Active Pharmaceutical Ingredients can be marked with a "Δ" if the product is sedative and a "R" if the product should be stored in a refrigerator

Fourth column describes under "Adm.form" whether the Active Pharmaceutical Ingredients should be available as capsules, tablets, suppositories injection liquid, infusion liquid, lotion, ointment, divisional powder, mouth spray, effervescent tablets, enema, oral solution or inhalation.

Fifth column describes the strength per tablet or per ml.

Sixth column describes the minimum amount the vessel must carry. The vessel may carry greater amounts than described.

Seventh column MFAG A2 describes the supplement the ship must bring if it sails with A-record AND has an evacuation time less than 24 hours.

Note: The columns marked MFAG describes the minimum quantities of additional (supplement) Active Pharmaceutical Ingredients the ship must bring, if the ship is carrying dangerous goods.

Besætningens medbragte medicin

Besætningen har ret til at medbringe egen medicin under den forudsætning, at de til stadighed har en gyldig helbredsundersøgelse for søfarende og fiskere. Medbragte lægemidler skal oplyses til skibsføreren eller sygdomsbehandleren.

Lægemidlet skal indskrives med følgende oplysninger:

- Lægemidlets navn
- Navn på besætningsmedlem
- Mængde og antal
- Daglig dosis
- Dato ind
- Eventuel kvittering for modtagelsen. (Kun hvis lægemidlet opbevares på sygelukafet)
- Dato ud
- Kvittering for modtagelsen (Kun hvis lægemidlet har været opbevaret på sygelukafet)

Guide til:

Fortegnelse og kontroldokument medicinsk udstyr

Publikationen indeholder en fortegnelse over det medicinske udstyr, der skal være om bord.

Rækkerne

Det medicinske udstyr er organiseret i hovedgrupper efter deres hovedanvendelsesområde. Eksempelvis er alle forbindinger og plastre organiseret i gruppe B.

Første kolonne beskriver det enkelte medicinske udstyrs bogstav og nummer. Eksempelvis har "Elastisk gazebind, ca. 4 cm x 4 m", kombinationen nummeret B.9, hvor bogstavet før punktum referer til hovedgruppen, og tallet efter referer til det specifikke medicinske udstyr.

Anden kolonne beskriver det specifikke medicinske udstyr.

Tredje kolonne beskriver den mindste mængde, skibet skal medbringe. Skibet må gerne medbringe større mængder end det beskrevne.

Fjerde kolonne. MFAG A2 beskriver det supplement, skibet skal medbringe, dersom det sejler med A-fortegnelse OG har en evakuerings tid på under 24 timer.

Bemærkning: Kolonnerne mærket MFAG beskriver de minimumsmængder af supplerende medicinsk udstyr som skibet skal medbringe, dersom skibet sejler med farligt gods.

Det mobile genoplivningsudstyr A.1

Søfartsstyrelsen har fastsat regler om ilt og det mobile genoplivningsudstyr i gældende Meddelelser A kapitel IX B. Man kan derudover pakke førstehjælpsudstyr og lægemidler til førstehjælp sammen med det mobile genoplivningsudstyr. Der er udarbejdet forslag til pakning af et mobilt genoplivningsudstyr på www.soefartsstyrelsen.dk.

Etiketter

Publikationen indeholde en kopi af de etiketter, der skal være på lægemiddelstofferne ved leveringen. Kopien kan anvendes som kontrol af, at lægemidlet er korrekt og påført den korrekte etiket. Er det ikke tilfældet, bør skibsføreren retlede sin leverandør.

Ved køb af lægemiddelstoffer, kan en kopi af etiketterne afleveres ved leverandøren. Derefter skal leverandøren påføre de korrekte etiketter, før udlevering

Private medicine of crewmembers

The crew has a right to bring their own medicine, provided they possess a valid health certificate for seafarers and fishermen at all time. The master or the person in charge of medical care has to be informed about medicine brought along.

The medicine must be registered with the following information:

- Name of the medicine
- Name of crewmember
- Amount
- Daily dosage
- Date in
- Possible receipt for received medicine.
(Only if the medicine is stored at the sickbay)
- Date out
- Sign when medicine is returned
(Only if the medicine has been stored in the sickbay)

Guide to:

Inventory and Control Document of Medical Equipment

The publication contains an inventory of the mandatory medical equipment onboard

The rows

The medical equipment is organized into main groups according to their main field of application. For example all dressings and plasters are organized in group B.

First column describes each individual medical equipment by number. The "elastic gauze", approx. 4 cm x 4 m, carries the combination B.9. The letters before the dot refers to the main group, and the number after the dot refers to the specific medical equipment.

Second column describes the specific medical equipment.

Third column describes the minimum amount the vessel must carry. The vessel may carry larger quantities than described.

Fourth column MFAG A2 describes the supplement the ship must bring if it sails with A-record has an evacuation time less than 24 hours.

Note: The columns marked MFAG describes the minimum quantities of additional (supplement) medical equipment the ship must bring, if the ship is carrying dangerous goods.

The Mobile Resuscitation Equipment A.1

The Danish Maritime Authority has established rules for oxygen and the mobile resuscitation equipment in the current Notice A chapter IX B. In addition you may pack equipment and medicine suited for first aid with the resuscitation equipment. Suggestions for the list may be found on www.dma.dk.

Labels

The publication contains copies of the labels that must be on each active pharmaceutical ingredient, at the time of delivery. The copy may be used as verification of proper product and label. If labelling is incorrect, the master should guide his supplier.

When purchasing Active Pharmaceutical Ingredients a copy of the label may be delivered to the supplier. Then the supplier applies the correct labels before delivery.

Indeks

Indekset henviser til lægemiddelstoffernes brugervejledning. I indekset kan du søge på visse sygdomme, symptomer og alle lægemiddelstoffer.

Rekvition

I publikationen findes en bestillingsliste, hvor du blot udfylder de ønskede mængder af lægemiddelstof og medicinsk udstyr.

Medicinkiste- / Mobilt genoplivningsudstyr certifikat.

Mindst en gang årligt skal skibets lægemiddelstoffer og medicinske udstyr gennemgås.

Det mobile genoplivnings udstyr skal kontrolleres hver 3 måned. Ved gennemgangen kontrolleres mængden, opbevaringsforhold, udløbsdato og etiketter. Nåle, kanyler og metalinstrumenter kontrolleres for rust. Kontroller, at emballagen er intakt. Vær opmærksom på, at lægemiddelstof og medicinsk udstyr skal opbevares tørt og mørkt, og ikke må opbevares i nærheden af klor.

Dokumentation af 3 måneders og årlig inspektion.

På side 1 og 2 er Medicinkiste – og mobilt genoplivningsudstyrs certifikat, som også kan anvendes til dokumentation af inspektionerne. Siderne udskrives og underskrives af skibsføreren.

Dispensation for andet lægemiddelstof

Det er ikke til alle tider, og i alle lande muligt at indkøbe det foretrukne lægemiddelstof. Den danske Søfartsstyrelse og Radio Medical Danmark har derfor udarbejdet en liste med godkendte alternativer. Hvis det er muligt at indkøbe lægemiddelstoffer fra listen med godkendte alternativer med den korrekte koncentration, og indgiftsmåde, er det ikke nødvendig at ansøge om dispensation.

Hvis apotek/forhandler tilbyder lægemiddelstof, som ikke findes på nogen liste, skal der søges om dispensation for anden medicin. Dette gøres på Søfartsstyrelsen hjemmeside, hvor der er en procedure for "Dispensation for anden medicin".

Listen over godkendte alternative, og dertil hørende labels og brugervejledninger kan downloades fra www.soefartsstyrelsen.dk.

Listen med godkendte alternativer findes i Excel format.

Hjælp

Er der noget du er i tvivl om, står Søfartsstyrelsens "Center for det Maritime Sundhedsvæsen" til din rådighed. Adressen er:

CMS, Vestervejen 1, 6720 Fanø
Telefon: +45 7219 6004
Email: cms@dma.dk

Index

The index refers to the user instruction of active pharmaceutical ingredients. You can search for diseases, symptoms and Active Pharmaceutical Ingredients in the index.

Requisition

In the publication is a requisition form available. You simply fill in the required quantities of the Active Pharmaceutical Ingredients and medical equipment.

Medicine chest- / Mobile resuscitation equipment certificate.

At least once a year the ship's Active Pharmaceutical Ingredients and medical equipment must be reviewed.

The mobile resuscitation equipment every 3 month.

When reviewing the quantities, storage conditions, expiry date and labels are controlled. Needles, cannulas and metal instruments are checked for rust. Check if the packaging is intact. Be aware that the Active Pharmaceutical Ingredients and equipment must be stored dry and dark, and not near chlorine.

Documentation of 3 months and annual inspection.

On page 1 and 2 is Medicine chest- and mobile resuscitation equipment's certificate's, it can be used to document the inspections. The page's has to be printed and the Master has to sign them.

Exemption for other active pharmaceutical ingredients

It is not at all times and in all countries possible to purchase the preferred Active Pharmaceutical Ingredients. The Danish Maritime Authority and Radio Medical Denmark have therefore drawn up a list of approved alternatives. If it is possible to purchase Active Pharmaceutical Ingredients from the list of approved alternatives with the correct concentration and method of administration, it is not necessary to apply for exemption.

If the pharmacy / retailer offer Active Pharmaceutical Ingredients that is not on the inventory- or alternative list, you must apply for an exemption. This is done on the DMA website, where there is a procedure for "exemption of medicine".

The list of approved alternative, and associated labels and user instructions can be downloaded from www.dma.dk. The list of approved alternatives is available in Excel format.

Help

If you have any questions, the Danish Maritime Authority's "Centre of Maritime Health Service" is at your disposal. The address is:

CMS, Vestervejen 1, 6720 Fanø
Telephone: +45 7219 6004
E-mail: cms@dma.dk

Anvendte forkortelser m.v. i Fortegnelse og Brugervejledning

afd. Pul.	=	afdelte pulvere
gel	=	gel til påsmøring på hud
htgl.	=	hætteglas
i. m.	=	intramuskulært = i musklen
inf. Væs.	=	infusionsvæske (til intravenøs brug)
inh. Pul.	=	inhalationspulver
inj. væske	=	injektionsvæske (til i.m. eller s.c.injektion)
i.v.	=	intravenøst = i en vene
"K"	=	opbevares i køleskab
klysma	=	klysma til anvendelse i endetarmen
linim.	=	liniment
gas.	=	luftformig gasart til medicinsk anvendelse
mixtur	=	væske som drikkes
næsedr.	=	næsedråber
opløsn.	=	opløsning
s.c.	=	subcutant = under huden
supp.	=	suppositorier = stikpille til brug i endetarmen
Δ	=	sløvende medicin, kan påvirke evnen til maskinbetjening og vagttjeneste
π	=	Receptpligtig medicin

Used abbreviations etc. In the Inventory and User instruction

d.p.	=	divisional powder
gel	=	gel for application on the skin
c.v	=	cappedvials
i.m.	=	intramuscular = in the muscle
inf. fl.	=	infusion fluid (for intravenous use)
inh. p.	=	inhalation powder
inj. subst.	=	injection substance (for reconstitution before use)
inj. liquid	=	injection fluid (for i.m. or s.c. injection)
i.v.	=	intravenous = in a vein
"R"	=	stored in refrigerator
enema	=	enema for use in the rectum
linim.	=	liniment
gas.	=	gas for medical use
mixture	=	drinkable liquid
nose d.	=	nose drops
diss.	=	dissolution
s.c.	=	subcutaneous = under the skin
supp.	=	suppositories = suppository for rectal use
Δ	=	sedative medicine, may effect the ability to operate machinery and bridge duty
π	=	Prescription drugs

Brugervejledning Lægemiddelstoffer

**User instruction Active Pharmaceutical
Ingredients**

Grp. 1 Ilt / Oxygen

Bevidstløs uden vejtrækning Bevidstløs med vejtrækning Medtaget almen tilstand Slag mod kraniet Større kvæstelser Vejtrækningsbesvær M FAG tabel 2, 3, 5, 9, 11	1.1+1.3	Lægemiddelstof:	Medicinsk ilt
		Form:	Luft på trykflasker.
		Virkning:	Tilfører ilt til cellerne via de røde blodlegemer.
		Dosering:	Bevidstløse uden vejtrækning: Manuel ventilation med 15 liter per minut. Bevidstløse med vejtrækning og øvrige tilstande: Maske med 1 til 9 liter/min.
		Bivirkninger:	Ilt er ufarligt til medicinsk brug.
		Holdbarhed:	Flaske: 3 år fra produktionsdato (Conoxia®, Linde Gas Therapeutics AGA).
		Opbevaring	Holdes væk fra brændbare gasser og stoffer. Flasker holdes under 50°C.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Unconscious without breathing Unconscious with breathing Weak general condition Blows to the skull Severe injuries Breathing difficulties M FAG table 2, 3, 5, 9, 11	1.1+1.3	Active Pharmaceutical ingredients:	Medical oxygen
		Form:	Gas in pressurized bottles.
		Effect:	Induces oxygen to the cells via the red blood cells.
		Dosage:	Unconscious without breathing: Manual ventilation, 15 litres per minute (A.1). Unconscious with breathing or other states: Oxygen mask, 9 litres per minute.
		Side-effects:	Oxygen is not dangerous when used for medicinal purposes.
		Validity:	Bottles 3 years from production date (Conoxia®, Linde Gas Therapeutics AGA).
		Storage:	Keep away from flammable gases and substances. Keep bottle under 50°C.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 2 Lokalbedøvelse / Local Anaesthetics

Lokalbedøvelse Syning af sår	2.1	Lægemiddelstof:	Lidocain
		Form:	Injektionsvæske 10 mg/ml til subcutan injektion.
		Virkning:	Lokalbedøver på injektionsstedet. Virker efter 1 - 2 minutter. Virker i 2 - 3 timer.
		Dosering:	Mængden afhænger af sårets størrelse. Normalt anvendes 20 - 50 mg. Maksimaldosis er 20 ml svarende til 200 mg. Se Søfartsstyrelsens Lægebog.
		Bivirkninger:	Allergi er sjælden. Ved utilsigtet injektion i blodbanen kan der opstå: 1. føleforstyrrelser, svimmelhed, uklarhed og øresusen. 2. synsforstyrrelser, muskelrysten og kramper. 3. blodtryksfald og langsom puls.
		Holdbarhed:	3 år fra produktionsdato (Xylocain®, AstraZeneca).
		Opbevaring:	Almindelig temperatur. Må ikke fryses.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Lokalbedøvelse skal ske i et rum med adgang til adrenalin (4.1) og genoplivningsudstyr (A.1). Risiko for alvorlige bivirkninger, hvis lidocain injiceres i blodbanen.

Local anaesthetics Suture of wounds	2.1	Active Pharmaceutical ingredients:	Lidocaine
		Form:	Injection liquid, 10 mg/ml for subcutaneous injection.
		Effect:	Anaesthetises the place of injection. Takes effect after 1-2 minutes. Effective for 2-3 hours.
		Dosage:	The amount depends on the size of the wound. Normally, 20-50 mg is used. The maximum dose is 20 ml, equivalent to 200 mg. See the Danish Maritime Authority's Medical Manual.
		Side-effects:	Allergy seldom occurs. In case of an unintended injection into the bloodstream, the following side-effects may occur: 1. affective disturbances, dizziness, confusion, and buzzing in the ears. 2. visual disturbances, muscle contractions, and convulsions. 3. falling blood pressure and slow pulse.
		Validity:	3 years from the date of production (Xylocain®, AstraZeneca).
		Storage:	Normal temperature. Not to be frozen.
		Remarks:	Consult <i>Radio Medical</i> . Local anaesthetics must be administered in a room with access to adrenaline (4.1) and resuscitation equipment (A.1). Risk of serious side-effects if lidocaine is injected directly into the bloodstream.
Lokalbedøvelse Kløestillende	2.2	Lægemiddelstof:	Lidocain
		Form:	Gel, 20 mg/g (2%).
		Virkning:	Lokalbedøver hud og sårrende. Kløestillende.
		Dosering:	Anvendes som lokalbedøvelse ved smertende sår eller smertefulde procedurer, fx ved rengøring af brandsår eller større og forurenede sår. Et tykt lag gel påsmøres huden, der herefter afdækkes lufttæt i ca. 15 minutter. Virker efter ca. 15 minutter. Virker i ca. 30 minutter.
		Bivirkninger:	Allergi er sjælden.
		Holdbarhed:	2 år fra produktionsdato (Xylocain®, AstraZeneca).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Local anaesthetics Relieves itching	2.2	Active Pharmaceutical ingredients:	Lidocaine
		Form:	Gel, 20 mg/g (2%).
		Effect:	Anaesthetises the skin and the edges of wounds. Relieves itching.
		Dosage:	To be used as local anaesthetics in case of painful wounds or procedures, e.g. cleansing of burns or large, dirty wounds. A thick layer is applied to the skin, which is subsequently covered in an air-proof condition for about 15 minutes. Takes effect after about 15 minutes. Effective for about 30 minutes.
		Side-effects:	Allergy seldom occurs.
		Validity:	2 years from the date of production (Xylocain®, AstraZeneca).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 3 Smertestillende midler / Analgetics

Feber Svage smerter: Hovedpine Muskelsmerter Ledsmarter Rygsmerter MFAG tabeller: 7, 8, 13	3.1	Lægemiddelstof:	Paracetamol
		Form:	Tabletter à 500 mg.
		Virkning:	Febernedsættende og smertestillende.
		Dosering:	Voksne: 1 g 3 - 4 gange dagligt. Højest 4 g per døgn.
		Bivirkninger:	Doser over 15 - 20 g indebærer risiko for leverskade, leversvigt og død.
		Holdbarhed:	5 år fra produktionsdato (Pamol Nycomed DK ApS).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Ingen.
Fever Weak pains: Headache Muscle pains Pain in joints Back pains MFAG tables: 7, 8, 13	3.1	Active Pharmaceutical ingredients:	Paracetamol
		Form:	Tablets of 500 mg.
		Effect:	Pain relieving and reduces fever.
		Dosage:	Adults: 1 g three-four times daily. Not more than 4 g daily.
		Side-effects:	Doses of more than 15-20 g involve liver damage, liver failure, and death.
		Validity:	5 years from the date of production (Pamol Nycomed DK ApS).
		Storage:	Normal temperature.
		<i>Remarks:</i>	None.
Moderate smerter: Menstruationssmerter Ledsmarter Knoglesmerter Knoglebrud Muskelsmerter	3.2	Lægemiddelstof:	Ibuprofen
		Form:	Tabletter à 400 mg.
		Virkning:	Smertestillende og febernedsættende.
		Dosering:	Voksne: 400 mg 3 - 4 gange dagligt. Højest 1800 mg per døgn. Undtagelsesvist op til 2400 mg per døgn i kortere perioder (4 - 6 uger).
		Bivirkninger:	Ondt i maven.
		Holdbarhed:	5 år fra produktionsdato (Ibumetin®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes ved: 1. Allergi for acetylsalicylsyre. 2. Astmapatienter. 3. Patienter i blodfortyndende behandling. 4. Patienter med symptomer på mavesår.

<p>Moderate pains: Period pains Pain in joints Bone-ache Fractures Muscle pains</p>	3.2	Active Pharmaceutical ingredients:	Ibuprofen
		Form:	Tablets of 400 mg.
		Effect:	Pain relieving and reduces fever.
		Dosage:	Adults: 400 mg three-four times daily. Not more than 1800 mg daily. Exceptionally, up to 2400 mg daily for shorter periods of time (4-6 weeks).
		Side-effects:	Stomach ache.
		Validity:	5 years from the date of production (Ibumetin®, Nycomed Danmark).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used in case of: 1. Allergy to acetylsalicylic acid. 2. Asthma patients. 3. Patients in blood diluting treatment. 4. Patients with symptoms of ulcer.
<p>Stærke smerter: Store sårskader Kvæstelser Forbrændinger Knoglebrud Nyresten Galdesten Blodprop i hjertet (se førstehjælp under 9.1 Nitroglycerin)</p> <p>MFAG tabel 7, 8,10,13</p>	3.4	Lægemiddelstof:	Morfin Δ
		Form:	Injektionsvæske 10 mg/ml, ampuller à 1 ml.
		Virkning:	Smertestillende. Vækker velvære, virker afslappende. Virker efter 5 – 15 min ved intramuskulær (i.m.) eller subcutan (sc.) injektion. Virker omgående ved intravenøs (i.v.) injektion. Virkningsvarighed: 3-4 timer.
		Dosering:	Subcutant og intramuskulært (voksne) 10 mg. Kan gentages, hvis tilstrækkelig effekt udebliver. Intravenøst (voksne): 5 mg (halv ampul) injiceres langsomt. Kan gentages, hvis tilstrækkelig effekt udebliver.
		Bivirkninger:	Svækker vejtrækningen. Kvalme og opkastning. Hæmmer optagelsen af medikamenter fra mavesækken. Kan give forstoppelse og hæmme blæretømningen.
		Holdbarhed:	2 år fra produktionsdato, (Morfin, Skanderborg apotek).
		Opbevaring:	Almindelig temperatur. Beskyttes mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved overdosering kan vejtrækningen ophøre. Modgift: Naloxone (5.1). Denne bivirkning og morfins virkning i øvrigt forstærkes af andre beroligende stoffer som alkohol, sovemidler og lignende. Overdosering er især en risiko ved hurtig injektion. Overdosering sker sjældent ved almindelig dosering. Observer vejtrækningen og ventilere om nødvendigt med ilt (1.1) på ventilationspose og maske (A.1).

Severe pain Major wounds Injuries Burns Fractures Kidney stone Gall stone Coronary thrombosis (see first aid under 9.1 Glyceryl trinitrate) MFAG Tabel 7,8,10,13	3.4	Active pharmaceutical ingredients:	Morphine Δ
		Form:	Injection liquid, 10 mg/ml, ampoules of 1 ml.
		Effect:	Pain relieving. Gives the patient a feeling of well-being, relaxant. Takes effect after about 5-15 minutes in case of intramuscular or subcutaneous injection. Takes effect immediately in case of intravenous injection Effective for 3-4 hours.
		Dosage:	Subcutaneously and intramuscularly (adults): 10 mg. May be repeated if the effect is not sufficient. Intravenously (adults): 5 mg (1/2 ampoule) to be injected slowly. May be repeated if the effect is not sufficient.
		Side-effects:	Restricts breathing. Nausea and vomiting. Restricts the absorption of medicaments from the stomach. May cause constipation and restrict bladder emptying.
		Validity:	2 years from the date of production (Morphine, Skanderborg Pharmacy).
		Storage:	Normal temperature. To be protected from light.
		Remarks:	Consult <i>Radio Medical</i> . In case of an overdose, the patient may stop breathing. Antidote: Naloxone (5.1). This side-effect and the effect of morphine as drowsiness may be increased by the use of other sedatives, such as alcohol, sleeping medicine, ect. Risk of overdose primarily if the injection is given too quickly. Overdoses are rare in case of normal dosage. Observe the breathing and if necessary, ventilate by means of oxygen (1.1) ventilations bag and mask (A.1).
Inflammation i huden Gigt Smerter	3.7	Lægemiddelstof:	Diclofenac
		Form:	Gel 11,6 mg/g.
		Virkning:	Smertestillende.
		Dosering:	Smøres på huden i et tyndt lag 3-4 gange i døgnet.
		Bivirkninger:	Rødme, hududslæt, hudkløe, eksem, hududslæt ved kontakt med midlet.
		Holdbarhed:	3 år, fra produktionsdato (Voltaren®gel, Novatis Healthcare
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må kun anvendes på intakt hud. Må ikke anvendes i ansigtet.
Inflammation of the skin. Arthritis Pain.	3.7	Active pharmaceutical ingredients:	Diclofenac
		Form:	Ointment with 11,6 mg/g.
		Effect:	Analgesic.
		Dosage:	Apply a thin layer to the skin 3-4 times daily.
		Side-effects:	Redness, rash, itching, eczema, skin rashes when in contact with the product.
		Validity:	3 years from the date of production (Voltaren®gel, Novatis Healthcare A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Only to be used on intact skin. May not be used in the face.

Nyresten Galdesten Ledsmerter	3.8	Lægemiddelstof:	Diclofenac
		Form:	Suppositorier 50 mg.
		Virkning:	Smertestillende.
		Dosering:	50 -100 mg mg x 1 dagligt.
		Bivirkninger:	Nedsat appetit, sure opstød og halsbrand, kvalme, opkastning, mavesmerter, diarré, luft i maven. Hovedpine, svimmelhed. Hududslæt og hudkløe.
		Holdbarhed:	3 år fra produktionsdato, (Voltaren, 2care4).
		Opbevaring:	Almindelig temperatur - ikke over 30°C.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes ved: 1. Allergi for acetylsalicylsyre. 2. Astmapatienter. 3. Patienter i blodfortyndende behandling. 4. Patienter med symptomer på mavesår.
Kidney stone Gall stone Pain in joint	3.8	Active pharmaceutical ingredients:	Diclofenac
		Form:	Suppositorie 50 mg.
		Effect:	Analgesic.
		Dosage:	50-100 mg ones daily.
		Side-effects:	Loss of appetite, heartburn and acid regurgitation, nausea and vomiting, abdominal pain, diarrhea, flatulence, headache, dizziness, skin rash, itchy skin.
		Validity:	3 years from the date of production (Voltaren,2care4).
		Storage:	Normal temperature - not above 30°C.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used in case of: 1. Allergy to acetylsalicylic acid. 2. Asthma patients. 3. Patients in blood diluting treatment. 4. Patients with symptoms of ulcer.
Stærke smerter	3.9	Lægemiddelstof	Morfin Δ
		Form:	Tablet 10 mg
		Virkning:	Smertestillende
		Dosering:	10 – 30 mg hver 4 time
		Bivirkninger:	Kvalme, forstoppelse, opkastningfeber, humørsvingninger og lavt blodtryk. Svimmelhed, hallucinationer, nedsat verjtrækning og mave smerter.
		Holdbarhed:	4 år fra produktionsdato (Morfin, EQL Pharma AB)
		Opbevaring:	Normal temperatur. Dog under 25 °C
		Anmærkninger:	Søg samråd med Radio Medical.

Severe pain	Active Pharmaceutical Ingredients:	Morphine Δ
	Form:	Tablets 10 mg
	Effect:	Pain relieving
	Dosage:	10 – 30 mg every 4 hours.
	Side-effects:	Nausea, constipation, vomiting, fever, mood swings and low blood pressure. Dizziness, hallucinations, decreased circulation and abdominal pain.
	Validity:	4 years from production date (Morphine, EQL Pharma AB)
	Storage:	Normale temperature – not above 25 °C
	Remarks:	Consult Radio Medical.

Grp. 4 Midler mod allergi, shock og søsyge / Agents for Allergy, Shock and Seasickness

Allergisk shock	4.1	Lægemiddelstof:	Adrenalin
		Form:	Injektionsvæske 1 mg/ml, ampuller à 1 ml til intramuskulær injektion.
		Virkning:	Øger pulsen, hjertets pumpekraft og blodtrykket. Sammenækning af blodkar. Virker 2 - 5 minutter efter intramuskulær injektion. Omsættes meget hurtigt. Halveringstiden i blodet er 2 - 3 minutter.
		Dosering:	Omgående (voksne): 0,6 mg (godt en halv ampul) dybt intramuskulært. Kan eventuelt gentages, hvis virkningen udebliver.
		Bivirkninger:	Blodtryksstigning, hurtig puls, rysten og svimmelhed.
		Holdbarhed:	2 år fra produktionsdato (Adrenalin®, 2care4),
		Opbevaring:	Læs indlægsseddel
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Er livsfarligt at anvende ved shock forårsaget af blod- eller væsketab.
Allergic shock	4.1	Active pharmaceutical ingredients:	Epinephrine
		Form:	Injection liquid, 1 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Increases the pulse, the pumping power of the heart, and the blood pressure. Contraction of blood vessels. Takes effect 2-5 minutes after intramuscular injection. Transformed very quickly. The half-life in the blood is 2-3 minutes.
		Dosage:	Immediately (adults): 0.6 mg (about half an ampoule) given deep intramuscular. May be repeated if the effect is insufficient.
		Side-effects:	High blood pressure, rapid pulse, shivering, and dizziness.
		Validity:	2 years from the date of production (Adrenalin®, 2care4)
		Storage:	Read packet insert
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Potentially fatal if used to treat shock caused by loss of blood or fluid.

Grp. 4 Midler mod allergi, shock og søsyge

(continued)

Allergi Høfeber Nældefeber Allergisk ansigtsødem Allergisk astma	4.2	Lægemiddelstof:	Acrivastin
		Form:	Kapsler à 8 mg.
		Virkning:	Modvirker allergi. Maksimal virkning efter 1½- 2 timer. Virkningsvarighed op til 8 timer.
		Dosering:	Voksne: 8 mg efter behov. Maksimalt 3 gange per døgn.
		Bivirkninger:	Døsighed kan forekomme og forstærkes af alkohol og beroligende midler, herunder sovemidler.
		Holdbarhed:	3 år fra produktionsdato (Benadryl®, McNeil ApS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Lægemidlet er ikke mærket med Δ men enkelte personer sløves alligevel så meget, at det er uforsvarligt med vagttjeneste eller maskinbetjening. Ved lungesympotomer: Se tillige gruppe 15.
Allergy Hay fever Urticaria Allergic face oedema Allergic asthma	4.2	Active pharmaceutical ingredients:	Acrivastine
		Form:	Capsule of 8 mg.
		Effect:	Eases allergic symptoms. Maximum effect after 1.5-2 hours. Effective for up to 8 hours.
		Dosage:	Adults: 8 mg when required. Maximum three times daily.
		Side-effects:	Drowsiness may occur and is increased by the use of alcohol and sedatives, including sleeping medicine.
		Validity:	3 years from the date of production (Benadryl®, McNeil Denmark ApS).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . The medicament is not marked with a Δ , but some patients experience a high degree of drowsiness, making bridge duty and operations of machinery unsafe. In case of symptoms of from the lungs: See group 15.
Allergi Allergisk astma Allergisk ansigtsødem	4.3	Lægemiddelstof:	Hydrocortison
		Form:	Injektionssubstans 100 mg i to-delt hætteglas med 2 ml sterilt vand til intramuskulært eller intravenøst brug.
		Virkning:	Modvirker allergi og inflammationer. Virkningen indtræder først 4 - 6 timer efter injektion, hvorfor anden akut behandling skal dække dette tidsrum.
		Dosering:	Voksne: 100 mg intramuskulært eller intravenøst som éngangsdosis.
		Bivirkninger:	Ved længere tids anvendelse: Forstyrrelser sukker, vand- og saltstofskiftet og i kirtlernes funktion. Sløring og/eller aktivering af infektioner. Eufori, sindssyge og søvnløshed. Ondt i maven og mavesår.
		Holdbarhed:	5 år fra produktionsdato (Solu-Cortef®, Pfizer ApS). Brugsfærdig injektionsvæske holdbart i 72 timer, hvis opbevaret i køleskab.
		Opbevaring:	Almindelig temperatur. Beskyttes mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved lungesympotomer: Se tillige gruppe 15.

Allergy Allergic asthma Allergic face oedema	4.3	Active pharmaceutical ingredients:	Hydrocortisonsuccinate
		Form:	Injection substance, 100 mg in capped vial divided into two sections, with 2 ml sterile water for intramuscular or intravenous use.
		Effect:	Eases symptoms of allergy and inflammations. Does not take effect until 4-6 hours after the injection. Other acute treatment to cover this period is needed
		Dosage:	Adults: 100 mg given intramuscularly or intravenously as a single dose.
		Side-effects:	If used for a extended period: Disturbances in the sugar, water and salt metabolism and in the glandula functions. Masking and/or activation of infections. Euphoria, mental illness and insomnia. Stomache ache and ulcer.
		Validity:	5 years from the date of production (Solu-Cortef®, Pfizer ApS). Injection liquid ready for use, valid for 72 hours, if stored cold.
		Storage:	Normal temperature. To be kept out of daylight. Injection liquid ready for use, to be stored refrigeated.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . In case of symptoms from lungs See also group 15.
Søsyge	4.5	Lægemiddelstof:	Cinnarizin Δ
		Form:	Tabletter à 25 mg.
		Virkning:	Modvirker søsyge.
		Dosering:	25 mg ½ time før effekt ønskes. Efter behov kan derefter tages 12,5 mg hver 6. time.
		Bivirkninger:	Døsighed.
		Holdbarhed:	5 år fra produktionsdato (Sepan®, Paranova Danmark A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Må ikke gives til patienter med grøn stær eller vandladningsbesvær (forstørret prostata).
Seasickness	4.5	Active pharmaceutical ingredients:	Cinnarizin Δ
		Form:	Tablets of 25 mg.
		Effect:	Prevent seasickness. Effect after 30 minutes.
		Dosage:	25 mg ½hour before desired effect. If needed, 12,5 mg may be given subsequently every 6 hours.
		Side-effects:	Drowsiness.
		Validity:	5 years from the date of production (Sepan®, Paranova Danmark A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Do not administer to patients with glaucoma or urinating problems (enlarged prostate).

Grp. 4 Midler mod allergi, shock og søsyge

(continued)

Allergiske reaktioner, hævelser osv. Oral behandling	4.6	Lægemiddelstof:	Prednisolon
		Form:	Tabletter á 25 mg.
		Virkning:	Bremser allergiske reaktioner, hævelser.
		Dosering:	Efter ordination af <i>Radio Medical</i> .
		Bivirkninger:	Forhøjet tryk i øjet, slører infektion. Infektion.
		Holdbarhed:	2 år fra produktions dato (Prednisolon, Actavis).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Allergic reactions, swellings etc. Oral treatment preferred.	4.6	Active Pharmaceutical ingredients:	Prenisolon
		Form:	Tablets of 25 mg.
		Effect:	Brakes allergic reation, swelling etc.
		Dosage:	After precipstion from <i>Radio Medical</i> .
		Side-effects:	Glaucoma, blurs infections, infektions.
		Validity:	2 years from productions date (Prednisolon, Actavis).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 5 Antidoter / Antidotes

<p>Modgift ved overdosering af morfin MFAG tabel 4, 13</p> <p>Førstehjælp: Ved udsættende vejtrækning understøttes manuelt med ilt på ballon og maske 15 l ilt/ minut.</p>	5.1	Lægemiddelstof:	Naloxon
		Form:	Injektionsvæske 0,4 mg/ml, ampuller à 1 ml til intramuskulær injektion.
		Virkning:	Naloxon ophæver morfins (bi)virkning. Virker 2 - 3 minutter efter intramuskulær injektion. Virkningsvarighed: Op til 4 timer.
		Dosering:	<ol style="list-style-type: none"> 1. Naloxon voksne: Der gives omgående 0,8 mg (2 ampuller) intramuskulært. 2. Hvis virkningen udebliver kan behandlingen gentages efter 5 minutter.
		Bivirkninger:	Rysten, utilpashed og opkastninger (abstinenssymptomer).
		Holdbarhed:	3 år fra produktionsdato (Naloxon, B. Braun).
		Opbevaring:	Beskyttes mod lys.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Morfin kan forringe eller ophæve vejtrækningen. Bivirkningen er sjælden og optræder sædvanligvis kun ved meget store overdoser og hyppigst i forbindelse med en hurtig indgivelse i venerne.
<p>Antidote to be administered in case of an overdose of morphine</p> <p>MFAG table 4, 13</p> <p>First Aid: If the patient stops breathing, assist breathing manually with oxygen 15l/minute on a mask.</p>	5.1	Active pharmaceutical ingredients:	Naloxone
		Form:	Injection liquid, 0.4 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Naloxone counteracts the (side)-effects of morphine. Takes effect 2-3 minutes after intramuscular injection. Effective for up to 4 hours.
		Dosage:	<ol style="list-style-type: none"> 1. Naloxone adults: 0.8 mg (2 ampoules) is given immediately intramuscular. 2. In case of no effect, the treatment may be repeated after 5 minutes.
		Side-effects:	Shivering, general indisposition, and vomiting.
		Validity:	3 years from the date of production (Naloxon B. Braun).
		Storage:	To be protected from daylight.
		Remarks:	Consult <i>Radio Medical</i> . Morphine may reduce or stop the patient's breathing. This side-effect is rare and, usually occurs in case of a large overdose and/or quick intravenous injection.

Insulintilfælde Insulinshock Lavt blodsukker	5.2	Lægemiddelstof:	Glucagon
		Form:	Ready-set bestående af: Injektionssubstans Glucagon 1 mg i hætteglas og sprøjte med 1 ml sterilt vand til intramuskulært brug.
		Virkning:	Frigør sukkerreserver fra leveren. Virker normalt inden 5 - 10 minutter. Virkningsvarighed: 10 - 40 minutter.
		Dosering:	Voksne: 1 mg intramuskulært. Kan evt. gentages efter 10 - 15 minutter. Når patienten er kommet til bevidsthed gives brød og sød saft.
		Bivirkninger:	Kvalme og opkastninger kan forekomme. Kan udløse hjertekrampe hos hjertesygge. Hjertekrampe behandles med glycerylnitrat spray (9.1).
		Holdbarhed:	Ved almindelig temperatur: 18 måneder fra produktionsdato (Glucagen® HypoKit, Novo Nordisk). Ved 2 - 8°C: 36 måneder fra produktionsdato. Den blandede opløsning skal anvendes umiddelbart.
		Opbevaring:	Almindelig temperatur eller 2 - 8°C. (Glucagen® HypoKit, Novo Nordisk).
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Dersom den blandede opløsning er uklar eller med krystaller, må blandingen ikke anvendes.
Hypoglycaemic coma Insulin shock Low blood sugar	5.2	Active pharmaceutical ingredients:	Glucagon
		Form:	Ready-set consisting of: Injection substance Glucagon, 1 mg in a capped vial and a syringe with 1 ml of sterile water for intramuscular use.
		Effect:	Releases sugar reserves from the liver. Normally, takes effect within 5-10 minutes. Effective for 10-40 minutes.
		Dosage:	Adults: 1 mg intramuscularly. May be repeated after 10-15 minutes. When the patient regains consciousness, he is given some bread and sweet juice.
		Side-effects:	Nausea and vomiting may occur. May cause angina pectoris in people with heart diseases. Angina pectoris is treated with glyceryl trinitrate spray (9.1).
		Validity:	At room temperature: 18 months from the date of production (Glucagen® HypoKit, Novo Nordisk). At 2-8°C: 36 months from the date of production. The mixed dissolution must be used immediately.
		Storage:	Normal temperature or 2-8°C. (Glucagen® HypoKit, Novo Nordisk).
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . If the mixed dissolution is cloudy or contains crystals, it must not be used.

MFAG tabel 17 Forgiftninger med insektgifte, der indeholder organophosphater eller carbamater.	5.3	Lægemiddelstof:	Atropin
		Form:	Injektionsvæske 1 mg/ml, ampuller à 1 ml til intravenøs injektion.
		Virkning:	Livreddende ved forgiftninger af visse insektgifte. Virkningen afhænger af gifttype og mængde. Se MFAG tabel 17.
		Dosering:	Dosis afhænger af forgiftningen. Atropin indgives langsomt intravenøst. Der startes med 2 mg (2 ampuller) og fortsættes med yderligere 2 mg efter hver 5 - 10 minutters mellemrum indtil pupillerne begynder at udvide sig. Giftig dosis er 10 - 20 mg til voksne. Dødelig dosis er ca. 200 mg til voksne.
		Bivirkninger:	Pupiludvidelse, lysskyhed, tørre slimhinder, hurtig puls, vandladningsbesvær og forstoppelse.
		Holdbarhed:	Plastampuller: 2 år fra produktionsdato. (Atropin®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
MFAG table 17 Poisoning with insecticides containing organophosphates or carbamates.	5.3	Active pharmaceutical ingredients:	Atropin
		Form:	Injection liquid, 1 mg/ml, ampoules of 1 ml for intravenous injection.
		Effect:	Life-saving in connection with poisonings caused by certain insecticides. The effect depends on the type and amount of poison. See MFAG table 17.
		Dosage:	The dose depends on the poisoning. Atropin is given slowly intravenously. At first, 2 mg (2 ampoules) is given, followed by another 2 mg after every 5-10 minutes until the pupils start dialating. Poisonous dose is 10-20 mg for adults. Lethal dose is about 200 mg for adults.
		Side-effects:	Dilation of the pupils, photophobia, dry mucous membranes, rapid pulse, dysuria and constipation.
		Validity:	Plastic ampoules: 2 years from the date of production (Atropin®, Nycomed Danmark).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
MFAG tabel 8, 16 Ætsninger med fluorholdige salte og syrer. Første hjælp: Huden skylles med rigelig vand i mindst 10 minutter. Forurenet tøj fjernes. Huden vaskes med vand og sæbe og skylles.	5.4	Lægemiddelstof:	Calciumgluconat 2%.
		Form:	Tuber à 25 g, indeholdende 2 g calciumgluconat per 100 g. Fremstilles af apoteket.
		Virkning:	Calcium neutraliserer i et vist omfang ætsningsskaderne ved fluoridætsninger.
		Dosering:	Huden indsmøres i calciumgluconat gel. Se vejledning i MFAG tabel 8 og 16.
		Bivirkninger:	Ingen.
		Holdbarhed:	2 år fra produktionsdato.
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .

<p>MFAG table 8, 16 Caustic burns caused by salts and acids containing fluorine.</p> <p>First aid: The skin is rinsed with plenty of water for at least 10 minutes. Contaminated clothes are removed. The skin is washed with water and soap and rinsed.</p>	5.4	Active pharmaceutical ingredients:	Calciumgluconate 2%
		Form:	Tubes of 25 g, containing 2 g of calciumgluconate per 100 g. Made by the individual pharmacy.
		Effect:	Calcium neutralizes caustic burns caused by fluorine to some extent.
		Dosage:	Calciumgluconate gel is applied to the skin. See guidelines in MFAG table 8 og 16.
		Side-effects:	None.
		Validity:	2 years from the date of production.
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
<p>MFAG tabel 16 Ætsninger med fluorholdige salte og syrer, der omfatter mere end 1% af hudens overflade.</p>	5.5	Lægemiddelstof:	Calciumlactogluconat og calciumcarbonat.
		Form:	Brusetabletter à 500 mg.
		Virkning:	Neutraliserer i et vist omfang skader efter optagelse af fluorsalte og fluorid forbindelser gennem huden.
		Dosering:	Ordination af <i>Radio Medical</i> .
		Bivirkninger:	Normalt ingen bivirkninger.
		Holdbarhed:	3 år fra produktionsdato (Calcium-Sandoz®, Novartis).
		Opbevaring:	Er fugtfølsomt. Opbevares tørt i tæt emballage ved almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
<p>MFAG table 16 Caustic burns caused by salts and acids containing fluorine and covering more than 1% of the surface of the skin.</p>	5.5	Active pharmaceutical ingredients:	Calciumlactogluconate and calciumcarbonate
		Form:	Effervescent tablets of 500 mg.
		Effect:	Neutralizes injuries caused by the absorption of fluoride compounds through the skin to some extent.
		Dosage:	Contact <i>Radio Medical</i> .
		Side-effects:	No side effects.
		Validity:	3 years from the date of production (Calcium-Sandoz, Orifarm).
		Storage:	Sensitive to humidity. To be stored in a dry place in a sealed packaging at room temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

MFAG tabel 10 Forgiftninger og ætsninger i mave og tarm.	5.6	Lægemiddelstof:	Aktivt kul
		Form:	50 g aktivt kul i granulatform til opblanding i 400 ml vand.
		Virkning:	Opsuger gift og lægemidler fra mavetarmkanalen og binder stofferne til kulpartiklerne.
		Dosering:	50 - 100 g kul (1 - 2 flasker) opslemmet i vand indtages så hurtigt som muligt. Ved alvorlige forgiftninger fortsættes med 20 g (160 ml) hver 6. time. Se også MFAG, Appendix 14.
		Bivirkninger:	Farver afføringen sort. Kan give forstoppelse. Nedsætter virkningen af lægemidler, der er indtaget via munden.
		Holdbarhed:	5 år fra produktionsdato (Carbomix Norit®). Efter opblanding i vand: Maks. 3 dage i køleskab.
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Anvendes ikke til bevidstløse. Må ikke anvendes ved indtagelse af stærke syrer/baser eller organiske opløsningsmidler (petroleumsignende).
MFAG table 10 Poisoning and caustic burns in stomach and intestines.	5.6	Active pharmaceutical ingredients:	Active carbon
		Form:	50 g of active carbon in the form of granulate for dilution in 400 ml of water.
		Effect:	Absorbs poison and medicaments from the gastrointestinal tract and binds the substances to the carbon particles.
		Dosage:	50-100 g carbon (1-2 bottles) suspended in water is taken as quickly as possible. In case of serious poisonings, continue with 20 g (160 ml) every sixth hour. See also MFAG, Appendix 14.
		Side-effects:	Black faeces. May cause constipation. Reduces the effect of medicaments taken through the mouth.
		Validity:	5 years from the date of production (Carbomix Norit®). After having been mixed with water, maximum 3 days in the fridge.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Do not use on unconscious patients. Not to be used in connection with strong acids/bases or organic solvents (kerosene related).

MFAG tabel 19 Forgiftninger med methanol og ethylene glycol	5.7	Lægemiddelstof:	Ethyl alkohol 99,5% (alternativt 99,9%)
		Form:	Væske
		Virkning:	Modgift mod methanol og ethylenglucols virkning.
		Dosering:	25 ml opblandes i 250 – 300 ml vand eller saft. Dosis gentages hver 3 time indtil evakuering. Se MFAG tabel 19.
		Bivirkninger:	Kvalme, hovedpine.
		Holdbarhed:	2 år fra produktionsdato.
		Opbevaring:	Almindelig stuetemperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
MFAG table 19 Poisoning caused by methanole and ethylene glycole	5.7	Active pharmaceutical ingredients:	Ethyl alcohol 99,5% (alternative 99,9%)
		Form:	Liquid.
		Effect:	Antidote against the effect of methanole and ethylene glucole.
		Dosage:	25 ml is mixed with 250-300 ml water or juice. Repeat every 3 hours until evacuation. See MFAG table 19.
		Side-effects:	Nausea, vomiting and headache.
		Validity:	2 years from the date of production.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 6 Midler mod psykiske lidelser og kramper / Mental disorder and Convulsion

Angst og urotilstande Alkoholabstinenser Forebyggelse af kramper Ledskred og brud 1. valgspræparat ved akut sindssyge (alternativt anvendes klysma (6.3))	6.2	Lægemiddelstof:	Diazepam Δ
		Form:	Tabletter à 5 mg.
		Virkning:	Stærkt sløvende og muskelafslappende. Dæmper angst, uro og rastløshed. Virker efter 30 minutter. Udskilles meget langsomt (Halveringstid: 20 - 50 timer). Forstærker virkningen af morfin, haloperidol, alkohol og andre sløvende medikamenter.
		Dosering:	5 mg x 3 dagligt. Akut sindssyge: Dosis efter aftale med <i>Radio Medical</i> . Se også 6.4.
		Bivirkninger:	Vanedannende. Døsighed, svimmelhed, forvirring, hukommelsessvigt og balanceforstyrrelser.
		Holdbarhed:	5 år fra produktionsdato (Diazepam, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Anxiety and states of restlessness Alcoholic withdrawal symptoms Prevention of convulsions Dislocations and fractures First choice of medicament in case of acute mental disease (alternatively, Enema (6.3) are used.)	6.2	Active pharmaceutical ingredients:	Diazepam Δ
		Form:	Tablets of 5 mg.
		Effect:	Strongly sedative and muscle-relaxant. Decreases anxiety and restlessness. Takes effect after 30 minutes. Is excreted very slowly (half-life: 20-50 hours). Increases the effect of morphine, haloperidole, alcohol and other sedatives.
		Dosage:	5 mg three times daily. Acute mental disease: Dosage after consultation with <i>Radio Medical</i> . See also 6.4.
		Side-effects:	Addictive. Drowsiness, dizziness, confusion, lapse of memory, and balance disturbances.
		Validity:	5 years from the date of production (Diazepam, Nycomed Danmark).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
Universelle kramper Kramper fra forgiftninger Abstinenskramper Epilepsi Akut sindssyge MFAG table 4, 5, 6	6.3	Lægemiddelstof:	Diazepam Δ
		Form:	Klysma, 4 mg/ml klysma til anvendelse i endetarmen.
		Virkning:	Stærkt sløvende og muskelafslappende. Dæmper angst, uro og rastløshed. Virker 5 - 10 minutter efter rektal administration. Udskilles meget langsomt (Halveringstid: 20 - 50 timer). Forstærker virkningen af morfin, haloperidol, alkohol og andre sløvende medikamenter.
		Dosering:	Efter aftale med <i>Radio Medical</i> - 1 klysma tømmes i endetarmen.
		Bivirkninger:	Døsighed, svimmelhed, forvirring, hukommelsessvigt og balanceforstyrrelser.
		Holdbarhed:	2,5 år fra produktionsdato (Stesolid®, Actavis A/S). Dog maksimalt 1 år efter anbrud af aluminiumsposen.
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .

Grp. 6 Midler mod psykiske lidelser og kramper / Mental disorder and Convulsion (continued)

Universal convulsions Convulsions caused by poisoning Withdrawal convulsions Epilepsy Acute mental illness MFAG tabel 4, 5, 6	6.3	Active pharmaceutical ingredients:	Diazepam Δ
		Form:	Enema, 4 mg /ml, enema for insertion into the rectum.
		Effect:	Strong sedative and muscle-relaxant. Decreases anxiety and restlessness. Takes effect 5-10 minutes after rectal administration. Is excreted very slowly (half-life: 20-50 hours). Increases the effect of morphine, haloperidole, alcohol and other sedatives.
		Dosage:	After consult of <i>Radio Medical</i> - 1 enema emptied into the rectum.
		Side-effects:	Drowsiness, dizziness, confusion, lapse of memory, and balance disturbances.
		Validity:	2,5 years from the date of production (Stesolid [®] , Actavis A/ S). However, maximum 1 year after having opened the aluminiumbag.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
2. valgspræparat ved akut sindssyge præget af mani, vildskab, psykisk og Motorisk uro, angst, vrangforestillinger og hallucinationer. Kvalme og opkastninger	6.4	Lægemiddelstof:	Haloperidol Δ
		Form:	Injektionsvæske 5 mg/ml, ampuller à 1 ml til intramuskulær injektion.
		Virkning:	Afdæmper symptomerne ved sindssygen. Dæmper kvalme og
		Dosering:	Akut sindssyge: 5 -10 mg intramuskulært. Kan eventuelt gentages 3 eller flere gange dagligt. Behandlingen kan eventuelt forstærkes med diazepam (6.2/ 6.3).
		Bivirkninger:	Den sløvende effekt forstærkes af alkohol, morfin og andre sløvende midler.
		Holdbarhed:	5 år fra produktionsdato (Serenase [®] , Janssen-Cilag).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Second choice of medicament in case of acute mental illness characterized by mania, wildness, mental and motoric restlessness, anxiety, delusions, and hallucinations. Nausea and vomiting	6.4	Active pharmaceutical ingredients:	Haloperidole Δ
		Form:	Injection liquid 5 mg/ml, ampoules of 1 ml for intramuscular injection.
		Effect:	Restricts the symptoms of mental disease. Reduces nausea and vomiting.
		Dosage:	Acute mental disease: 5-10 mg intramuscularly. May be repeated three times daily or more. The treatment may be enhanced by the use of diazepam (6.2/6.3).
		Side-effects:	The sedative effect is increased by alcohol, morphine, and other sedatives.
		Validity:	5 years from the date of production (Serenase [®] , Janssen-Cilag).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 6 Midler mod psykiske lidelser og kramper / Mental disorder and Convulsion (continued)

Søvnløshed Indsovningsbesvær	6.5	Lægemiddelstof:	Zolpidem Δ
		Form:	Tabletter à 10 mg.
		Virkning:	Stærkt sløvende. Virker hurtigt efter indtagelse. Virkningen varer op til 6 timer. Virkningen forstærkes af andre sløvende stoffer, herunder alkohol.
		Dosering:	Voksne 10 mg 15-30 min. før sengetid.
		Bivirkninger:	Kvalme, opkastning, diarré. Hovedpine, koncentrationsbesvær, konfusion, hukommelsesbesvær, svimmelhed, døsigthed, hudreaktioner, dobbeltsyn.
		Holdbarhed:	5 år fra produktionsdato (Zolpidem Hexal A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Insomnia Difficulty falling asleep	6.5	Active pharmaceutical ingredients:	Zolpidem Δ
		Form:	Tablets of 10 mg.
		Effect:	Strong sedative, rapid effect after ingestion. Effective up till 6 hours. The effect is increased by the use of other sedatives, including alcohol.
		Dosage:	Adults 10 mg 15-30 minutes. before bedtime.
		Side-effects:	Nausea, vomiting, diarrhea. Headache, difficulty concentrating, confusion, memory loss, dizziness, drowsiness, skin reactions, double vision.
		Validity:	5 years from the date of production (Zolpidem Hexal A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 7 Midler mod infektioner / Infections			
Halsbetændelse Mundhuleinfektioner Bihulebetændelse Mellemørebetændelse Lungebetændelse	7.1	Lægemiddelstof:	Phenoxyethylpenicillin
		Form:	Tabletter à 660 - 800 mg (ca.1 mill. IE).
		Virkning:	Dræber de hyppigst forekommende bakterier, der forårsager infektioner i luftvejene.
		Dosering:	400–800 mg × 2 – 4 dagligt.
		Bivirkninger:	Kvalme, diarré og hududslæt. Sjældent ses overfølsomhedsshock.
		Holdbarhed:	3 år fra produktionsdato (Primcillin®, MEDA).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Overfølsomhedsshock optræder meget sjældent, og som regel senest 30 minutter efter indgift. Behandling: Adrenalin 0.6 mg intramuskulært (4.1).
Inflammation of the throat Infections of the oral cavity Sinusitis Inflammation of the middle ear Pneumonia	7.1	Active pharmaceutical ingredients:	Phenoxyethylpenicillin
		Form:	Tablets of 660 - 800 mg (approx. 1 MIU).
		Effect:	Kills the most common bacteria causing infections of the respiratory tract.
		Dosage:	400-800 mg × 2-4 times daily.
		Side-effects:	Vomiting, diarrhoea and rash. Rarely allergic shock.
		Validity:	3 years from the date of production (Primcillin®, MEDA).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Allergic shock is very rare and usually occurs within 30 minutes after administration. Treatment: Adrenaline 0.6mg intramuscular (4.1).
Lungebetændelse efter inhalation af giftige kemiske stoffer MFAG tabel 9	7.2	Lægemiddelstof:	Amoxicillin
		Form:	Kapsler eller tabletter à 500 mg.
		Virkning:	Modvirker inflammation.
		Dosering:	375-750 mg 2 gange daglig.
		Bivirkninger:	Feber, kvalme, opkastning, sure opstød, diarré, hudkløe, hududslæt.
		Holdbarhed:	4 år fra produktionsdato (Imadrax, Sandoz A/S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Overfølsomhedsshock optræder meget sjældent og som regel senest 30 minutter efter indgift. Behandling: Adrenalin 0,6 mg intramuskulært.(4.1)
Pneumonia after inhalation of poisonous chemical substances MFAG tabel 9	7.2	Active Pharmaceutical Ingredients:	Amoxicilline
		Form:	Capula or tablets of 500 mg.
		Effect:	Prevent inflammation.
		Dosage:	375-750 mg twice daily.
		Side-effects:	Fever, nausea, vomiting, heartburn, diarrhoea, rash and urticaria, Rarely allergic shock.
		Validity:	4 years from the date of production (Imadrax, Sandoz A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Allergic shock is very rare and usually it occurs within 30 minutes after administration. Treatment: Adrenaline 0.6 mg intramuscular (4.1).

Grp. 7 Midler mod infektioner / Infections

(continued)

Halsbyld Tandbyld Knogleinfektion Underlivsinfektioner Tarminfektioner Blindtarmsbetændelse (tjek urinstiks G.1)	7.6	Lægemiddelstof:	Metronidazol
		Form:	Tabletter à 500 mg.
		Virkning:	Bakteriedræbende.
		Dosering:	500 mg x 3 dagligt.
		Bivirkninger:	Mavekneb, smagsforstyrrelser, hovedpine, hududslet og
		Holdbarhed:	5 år fra produktionsdato (Metronidazol, Actavis).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Boil in the throat Tooth abscess Bone infection Infection of female reproductions organs Abdominal infections Intestinal infections Appendicitis (Check dipstick G.1)	7.6	Active pharmaceutical ingredients:	Metronidazole
		Form:	Tablets of 500 mg.
		Effect:	Kills bacteria.
		Dosage:	500 mg three times daily.
		Side-effects:	Bellyache, altered sense of taste, headache, rash, and dizziness. The urine may be coloured black. May
		Validity:	5 years from the date of production (Metronidazol, Actavis).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
Halsbyld Tandbyld Knogleinfektion Underlivsinfektioner Tarminfektioner Blindtarmsbetændelse MFAG tabel 10	7.7	Lægemiddelstof:	Metronidazol
		Form:	Stikpille à 500 mg.
		Virkning:	Bakteriedræbende.
		Dosering:	500 mg x 3 daglig.
		Bivirkninger:	Kvalme, opkastning, mavesmerter, diarré, forstoppelse, og smagsforstyrrelse. Mørkfarvet urin. Kan virke som antabus ved samtidig indtagelse af alkohol.
		Holdbarhed:	3 år fra produktionsdato (Flagyl®, Sanofi-aventis)
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan om nødvendigt anvendes, til gravide. Anvendes ofte i kombination med andre antibiotika, fx Ciprofloxacin (7.8) eller Ceftriaxon (7.9). Må ikke anvendes sammen med 15.1 Dexofan.

Grp. 7 Midler mod infektioner / Infections		(continued)	
Boil in the throat Tooth abscess Bone infection Infection of female reproductions organs Abdominal infections Intestinal infections Appendicitis (Check dipstick G.1)	7.7	Active pharmaceutical ingredients:	Metronidazol
		Form:	Suppository of 500 mg.
		Effect:	Kills bacteria.
		Dosage:	500 mg three times daily.
		Side-effects:	Nausea vomiting, abdominal pain, diarrhea, constipation, and taste disturbance. Dark colored urine. May cause a reaction similar to Antabuse in connection with alcohol.
		Validity:	3 years from productions date (Flagyl®, Sanofi-aventis)
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . May be used during pregnancy, if nessasary. Often used in combination with other antibiotic,s e.g. Ciprofloxacin (7.8) or Ceftriaxon (7.9). May not be used with 15.1 Dextromethorphan.
Tandbyld Halsbyld Svær bronkitis Lungepest Akut diarré Kolera Dysenteri Blindtarmsbetændelse Gonoré Urinrørsbetændelse Nyrebækkenbetændelse (tjek urinstiks G.1)	7.8	Lægemiddelstof:	Ciprofloxacin
		Form:	Tabletter à 250 mg.
		Virkning:	Dræber bakterier.
		Dosering:	250 - 750 mg x 2 dagligt.
		Bivirkninger:	Mavekneb, kvalme, opkastninger, madlede, hovedpine, svimmelhed, træthed og kramper, muskel-, led- og seneømhed, allergi, overfølsomhedsreaktion ved samtidig
		Holdbarhed:	5 år fra produktionsdato (Cifin, PharmaCoDane ApS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Gives ved mange infektioner i kombination med metronidazol (7.6 eller 7.7). Må ikke gives til gravide eller ammende.

Grp. 7 Midler mod infektioner / Infections (continued)

Boil in the throat Tooth abscess Severe bronchitis Pneumonic plague Acute diarrhoea Cholera Dysentery Appendicitis Gonorrhoea Urethritis Pyelitis (Check dipstick G.1)	7.8	Active pharmaceutical ingredients:	Ciprofloxacin
		Form:	Tablets of 250 mg.
		Effect:	Kills bacteria.
		Dosage:	250-750 mg twice daily The tablets are given 1 hour before a
		Side-effects:	Bellyache, nausea, vomiting, loss of appetite. Headache, dizziness, fatigue, and convulsions. Soreness in the muscles, joints and tendons. Allergy. Allergic reaction in case of simultaneous exposure to
		Validity:	5 years from the date of production (Cifin, PharmaCoDane Aps).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Is given in combination with metronidazole (7.6 or 7.7) in connection with many infections. Should not be given to pregnant or breast-feeding patients.
		Blodforgiftning Sepsis Meningitis Lungebetændelse Blindtarmsbetændelse Nyrebækkenbetændelse Store sårskader Åbne knoglebrud MFAG tabel 10 (Ceftriaxon erstatter Cefuroxime)	7.9
Form:	Injektionssubstans, 1 g tørstof i hætteglas + 10 ml sterilt		
Virkning:	Bakteriedræbende.		
Dosering:	1 - 2 g intravenøst i døgnet i én dosis.		
Bivirkninger:	Allergiske reaktioner: Hududslet, nældefeber, ledsmerter, overfølsomhedsshock.		
Holdbarhed:	Injektionssubstans: 3 år fra produktionsdato (Rocephalin®, Roche).		
Opbevaring:	Må ikke opbevares over 30°C.		
<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Gives ved mange infektioner i kombination med metronidazol (7.6 eller 7.7). Er stærkt vævsirriterende ved intramuskulær injektion.		
Sepsis Blood poisoning Meningitis Pneumonia Appendicitis Pyelitis Major wounds Open fractures MFAG table 10 (Ceftriaxone replaces Cefuroxime)	7.9	Active pharmaceutical	Ceftriaxone
		Form:	Injection substance, 1 g of dry matter in a capped vial + 10 ml of sterile water in a capped vial for intravenous use. Can be given intramuscular - Contact <i>Radio Medical</i> .
		Effect:	Kills bacteria.
		Dosage:	1-2 g intravenously daily as a single dose.
		Side-effects:	Allergic reactions: Rash, urticaria, arthralgia, allergic shock.
		Validity:	Injection substance: 3 years from the date of production (Rocephaline®, Roche). Dissolution ready for use: 24 hours, if refrigerated.
		Storage:	Not to be stored at more than 30°C.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Often used in combination with metronidazole (7.6 or 7.7) in connection with many infections. In case of intramuscular injection, strong irritation of tissue.

<p>Stivkrampevaccine</p> <p>Gives ved ulykker med kvæstelser og sår, med mindre den skadede med sikkerhed er dækket med stivkrampe-vaccine.</p> <p>Vaccinen suppleres med tetanus immunglobulin (se 7.11).</p> <p>Vaccination skal ske i behandlingsrum med adgang til Adrenalin(4.1) og Genoplivningsudstyr (A.1).</p>	7.10	Lægemiddelstof:	Tetanus holdig vaccine (tetanustoxoid)
		Form:	Injektionsvæske
		Virkning:	Aktiverer patientens immunsystem.
		Dosering:	Følg ordination fra <i>Radio Medical</i> .
		Bivirkninger:	Lokal rødme, hævelse og ømhed. Hovedpine, utilpashed og feber ses undertiden og fortager sig efter et døgn. Ledsmerter og hævede, ømme lymfeknuder kan ses på 8. dag. Allergisk shock forekommer sjældent.
		Holdbarhed:	2 år fra produktionsdato (Tetanusvaccine SSI).
		Opbevaring:	Opbevares køligt mellem 2 - 8° C. Frossen vaccine må ikke anvendes.
	<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .	
<p>Tetanus vaccine</p> <p>To be administered in case of accidents causing injury and wounds unless the wounded person is for certain covered by a tetanus vaccine.</p> <p>The vaccine is to be supplemented by tetanus immune globulin (see 7.11).</p> <p>The vaccination must be given in a room with access to Epinephrine (4.1) and Resuscitation equipment (A.1).</p>	7.10	Active pharmaceutical ingredients:	Vaccine containing Tetanus (tetanustoxoid)
		Form:	Injection liquid.
		Effect:	Activates the patient's immune system.
		Dosage:	Follow prescription from <i>Radio Medical</i> .
		Side-effects:	Local blushing, swelling, and soreness. Headache, weak general condition, and fever are sometimes seen, which decrease after about 1 day. Arthralgia and swollen, sore lymph nodes may be seen on the eighth day.
		Validity:	2 years from the date of production (Tetanus vaccine SSI).
		Storage:	To be stored cool at between 2-8°C. Frozen vaccine must not be used.
	<i>Remarks:</i>	Consult <i>Radio Medical</i> .	

Grp. 7 Midler mod infektioner / Infections		(continued)	
Infektioner forårsaget af stafylokokker. Knogleinfektion	7.12	Lægemiddelstof:	Flucloxacillin
		Form:	Tabletter à 500 mg.
		Virkning:	Bakteriedræbende.
		Dosering:	750-1000 mg 3-4 gange dagligt.
		Bivirkninger:	Kvalme, opkastning, diarré, hududslæt.
		Holdbarhed:	2 år fra produktionsdato. Åbne blisterpakninger 6 måneder (Heracillin, Meda AS).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Infections caused by staphylococci. Infections in the bones	7.12	Active pharmaceutical ingredients:	Flucloxacillin
		Form:	Tablets of 500 mg.
		Effect:	Kills bacteria.
		Dosage:	750-1000 mg 3-4 times daily.
		Side-effects:	Nausea, vomiting, diarrhea, skin rash.
		Validity:	2 years from the date of production. 6 months for open blister pack(Heracillin, Meda AS).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .
Halsbetændelse Mundhuleinfektioner Bihulebetændelse Mellemørebetændelse Lungebetændelse Legionella Chlamydia infektioner Mavesårsinfektioner Alternativ til penicillin ved allergi MFAG tabel 9	7.15	Lægemiddelstof:	Azithromycin
		Form:	Tabletter á 500mg.
		Virkning:	Standser bakteriernes vækst.
		Dosering:	Voksne: 500 mg x 1-2 dagligt.
		Bivirkninger:	Mavesmerter, kvalme, nedsat appetit, kraftsløshed, svimmelhed, smagsforstyrrelser, hovedpine, hudkløe og hududslæt.
		Holdbarhed:	5 år fra produktionsdato (Azithromycin, KRKA).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Inflammation of the throat Infections of the oral cavity Sinusitis Inflammation of the middle ear Pneumonia Legionella Chlamydia infections Infections in connection with ulcer Alternative to penicillin in case of allergy MFAG table 9	7.15	Active Pharmaceutical ingredients:	Azithromycin
		Form:	Tablets of 500mg.
		Effect:	Stops bacterial growth.
		Dosage:	Adults: 500 mg x 1-2 daily.
		Side-effects:	Abdominal pain, nausea, decreased appetite, weakness, dizziness, taste disturbances, headache, itchy skin and skin rash.
		Validity:	5 years from date of production (Azithromycin, KRKA).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 7 Midler mod infektioner / Infections **(continued)**

Urinvejsinfektioner	7.16	Lægemiddelstof:	Pivmecillinam
		Form:	Tablet 400 mg
		Virkning:	Bakteriedræbende
		Dosering:	400 mg x 3-4 gange dagligt
		Bivirkninger:	Kvalme,diarre, træthed, nældefeber, hudkløe, hududslæt, mundsår, hovedpine.
		Holdbarhed:	3 år fra produktions dato (Pivmecillinamhydrochlorid,"Paranova")
		Opbevaring:	Almindelig temperatur
		<i>Anmærkninger:</i>	Søg samråd med Radio Medical
Urinary tract infection	7.16	Active Pharmaceutical Ingredients:	Pivmecillinam
		Form:	Tablets 400 mg
		Effect:	Kills bacteria
		Dosage:	400 mg x 3-4 times daily
		Side-effects:	Nausea, diarrhoea, fatigue, hives, skin itching, skin rash, mouth ulcers, headache.
		Validity:	3 years from production date (Pivmecillinamhydrochlorid "Paranova")
		Storage:	Normal temperature
		<i>Remarks:</i>	Consult Radio Medical
Herpes simplex Skoldkopper Herpes zoster (helvedesild)	7.17	Lægemiddelstof:	Aciclovir
		Form:	Tablet 400 mg
		Virkning:	Hæmmer virus produktionen
		Dosering:	400-800 mg x 3 times daily
		Bivirkninger:	Diarré, kvalme, opkast, feber, kraftsløshed, træthed, hovedpine, døsigthed, hudkløe, hududslæt, lysfølsom hud.
		Holdbarhed:	5 år fra produktions dato (Aciclovir, STADA Nordic ApS)
		Opbevaring:	Opbevares under 25 °C
		<i>Anmærkninger:</i>	Søg samråd med Radio Medical
Herpes simplex Variceller Herpes zoster	7.17	Active Pharmaceutical Ingredients:	Aciclovir
		Form:	Tablets 400 mg
		Effect:	Inhibits virus production
		Dosage:	400-800 mg x 3 times daily
		Side-effects:	Diarrhoea, nausea, vomiting, fever, weakness, fatigue, headache, drowsiness, itchy skin, skin rash, photosensitive skin.
		Validity:	5 years from productions date (Aciclovir, STADA Nordic ApS)
		Storage:	Do not store above 25°C
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 8 Malaria – forebyggelse / Malaria – Prevention

Forebyggelse af malaria	8.1	Lægemiddelstof:	Atovaquon/Proguanil (Malarone®)
		Form:	Tabletter à 250/100 mg.
		Virkning:	Forebygger malaria.
		Dosering:	Tabletterne <i>skal</i> indtages med fedtholdigt mad (mælk/smør). Forebyggelse fra 1 dag før ankomst til 7 dage efter afrejse fra malariaområdet. Dosis voksne: 1 tablet dagligt.
		Bivirkninger:	Hovedpine, mavekneb, kvalme, opkastning, appetitløshed, diarré, hoste og allergisk hududslet. Ved opkastning inden én time efter indtagelse af medikamentet indtages ny tablet.
		Holdbarhed:	5 år fra produktionsdato (Malarone®, GlaxoSmithKline Pharma A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Forebyggelse: Se Søfartsstyrelsens gældende vejledning om Malaria.
Prevention of malaria	8.1	Active pharmaceutical ingredients:	Atovaquone/Proguanil (Malarone®)
		Form:	Tablets of 250/100 mg.
		Effect:	Prevents malaria.
		Dosage:	The tablets <i>must</i> be taken with fatty food (milk/butter). The prevention starts 1 day before arrival and lasts until 7 days after departure from the malaria area. Dose, adults: 1 tablet a day.
		Side-effects:	Headache, stomach pains, nausea, vomiting, loss of appetite, diarrhoea, coughing and allergic rash. If the patient vomits within 1 hour after have taken the medicament, a new tablet has to be consumed.
		Validity:	5 years from the date of production (Malarone®, GlaxoSmithKline Pharma A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Primary prevention of malaria: See the Danish Maritime Authority's guide on Malaria.
Forebyggelse af malaria Anvendes til forebyggelse, dersom man ikke tåler 8.1	8.2	Lægemiddelstof:	Doxycyklin
		Form:	Tabletter à 100 mg.
		Virkning:	Forebygger malaria.
		Dosering:	Indtages med et glas vand eller et let måltid uden mælkeprodukter. Behandlingen udstrækkes fra 1 dag før ankomst til 4 uger efter afrejse fra malariaområdet. Dosis voksne og børn over 12 år: 1 tablet dagligt.
		Bivirkninger:	Kvalme, vaginal svamp og toksiske hudreaktioner i forbindelse med sollys.
		Holdbarhed:	5 år fra produktionsdato (Vibradox®, Sandoz).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes af gravide eller ammende eller af børn under 12 år. Direkte sollys bør undgås. Interaktion med mange lægemidler.

Grp. 8 Malaria – forebyggelse / Malaria – Prevention

(continued)

Prevention of malaria To be used for prevention if 8.1 is not tolerated	8.2	Active pharmaceutical ingredients:	Doxycycline
		Form:	Tablets of 100 mg.
		Effect:	Prevents malaria.
		Dosage:	To be taken with a glass of water or a light meal without any milk products. The prevention starts 1 day before arrival and lasts until 4 weeks after departure from the malaria area. Dose, adult and children over the age of 12 or 45 kg: 1 tablet daily.
		Side-effects:	Nausea, vaginal candidiasis, toxic skin reaction in connection with sunlight.
		Validity:	5 years from the date of production (vibradox®, Sandoz).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used by pregnant or breast-feeding woman or children below the age of 12. Direct sunlight should be avoided. Interaction with many other medicines.

Grp. 8 Malaria – behandling / Malaria – Treatment

Til behandling af P. Falciparum malaria. På baggrund af: Positiv malaria test for P. Falciparum eller almen sygdomsfornemmelse og feber kombineret med relevant risiko for malaria	8.4	Lægemiddelstof:	Artemether og lumefantrine
		Form:	Tabletter á 20mg/120mg.
		Virkning:	Dræbermalaria parasitterne.
		Dosering:	Efter kontakt med <i>Radio Medical</i> . Start omgående med 4 tbl. Derefter: 4 tabl 8 timer efter første dosis 4 tabl 24 timer efter første dosis. 4 tbl. 36 timer efter første dosis. 4 tbl. 48 timer efter første dosis. 4 tbl. 60 timer efter første dosis.
		Bivirkninger:	Mave-Tarm gener, hudkløe, hududslæt, hovedpine og svimmelhed.
		Holdbarhed:	2 år fra produktionsdato (Riamet®, Novatis).
		Opbevaring:	Må ikke opbevares over 30°C.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> uanset resultatet af malariatest.
Treatment of P. Falciperum malaria Based on: Positive Malaria Test for P. Falciparum or general malaise and fever combined with relevant risk of malaria	8.4	Active Pharmaceutical ingredients:	Artemether and lumefantrine
		Form:	Tablets of 20mg/120mg.
		Effect:	Kills the malaria parasite.
		Dosage:	Consult <i>Radio Medical</i> . 4 tablets as soon as possible. Then: 4 tablets 8 hours later, then 4 tablets 24 hours after the first dose, then 4 tablets 36 hours after the first dose, then 4 tablets 48 hours after the first dose and then 4 tablets 60 hours after the first dose.
		Side-effects:	Gastrointestinal disorders, itchy skin, skin rash, headache and dizziness.
		Validity:	2 years from production date (Riamet®, Novatis).
		Storage:	Do not store above 30°C.
		Remarks:	Consult <i>Radio Medical</i> regardless of the result of the malaria test.

Grp. 9 Hjerte og kredsløb / Cardiac and circuitry

<p>Hjertekrampe Smerter i hjertet Blodprop i hjertet</p> <p>Førstehjælp: Hjertestilling (elevator overkrop) 2 - 3 l ilt/minut Glycerylnitrat (9.1) Acetylsalicylsyre (9.3)</p> <p>Evt. for smerter: Morfin (3.4) / Tramadol (3.5)</p>	9.1	Lægemiddelstof:	Glycerylnitrat
		Form:	Mundhule spray med 0,4 mg/dosis.
		Virkning:	Udvider blodkar. Virker inden 1 minut. Virkningsvarighed: 10 - 30 minutter.
		Dosering:	Medikamentet pustes ind under tungen eller på kindens inderside. 1 pust så hurtigt som muligt efter smerter opståen. Kan gentages 1 gang. Hvis smerterne ikke fortager sig, suppleres med morfin (3.4).
		Bivirkninger:	Ansigtserødem, hovedpine og kvalme. Kan sænke blodtrykket lidt.
		Holdbarhed:	2 år fra produktionsdato (Glytrin®, Meda).
		Opbevaring:	Almindelig temperatur. Beskyttes mod frost, direkte varme og sollys.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Overvåg blodtrykket, som helst ikke må komme under 90 mmHg systolisk. Må ikke anvendes i nærheden af åben ild eller gløder.
<p>Angina pectoris Heart pain Coronary thrombosis</p> <p>First aid: Sit up in comfortable position 2-3 l oxygen/minute Glyceryl trinitrate (9.1) Acetylsalicylic acid (9.3)</p> <p>Possibly against pain: Morfin (3.4) / Tramadol (3.5)</p>	9.1	Active pharmaceutical ingredients:	Glyceryl trinitrate.
		Form:	Oral cavity spray with 0.4 mg/dose.
		Effect:	Expands the blood vessels. Takes effect within 1 minute. Effective for 10-30 minutes.
		Dosage:	The medicament is sprayed under the tongue or at the inner side of the cheek. 1 spray as soon as possible after the pain has occurred. May be repeated once. If the pain does not decrease, supplement with morphine (3.4).
		Side-effects:	Blushing, headache, and nausea. May cause a minor fall in blood pressure.
		Validity:	2 years from the date of production (Glytrin®, Meda).
		Storage:	Not to be stored above 25°C. Protect from frost, direct heat and sunlight.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Control the blood pressure, which should not be below 90 mmHg systolic. Not to be used close to open fire or embers.

Hjertekrampe Smerter i hjertet Blodprop i hjertet Blodfortynding Se også 9.1	9.3	Lægemiddelstof:	Acetylsalicylsyre
		Form:	Tabletter à 150 mg.
		Virkning:	Virker blodfortyndende. Forhindrer blodpropper.
		Dosering:	Akut: 300 mg som tygges. Vedligehold: 75 mg dagligt.
		Bivirkninger:	Øget blødningsrisiko. Risiko for allergiske reaktioner. Farlig dosis: 150 mg per kg legemsvægt.
		Holdbarhed:	5 år fra produktionsdato (Hjertemagnyl®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Bør ikke gives til patienter med mavesår.
Cardiospasm Heart pain Coronary thrombosis Blood-thinning See also 9.1	9.3	Active pharmaceutical ingredients:	Acetylsalicylic acid
		Form:	Tablets of 150 mg.
		Effect:	Thins the blood. Prevents thrombosis.
		Dosage:	Acute: 300 mg has to be chewed. Hereafter: 75 mg, daily.
		Side-effects:	Increased risk of bleeding. Risk of allergic reactions. Dangerous dosage: 150 mg per kg bodyweight.
		Validity:	5 years from the date of production (Hjertemagnyl®, Nycomed Denmark).
		Storage:	Normal temperature.
<i>Remarks:</i>	Consult <i>Radio Medical</i> . Not to be given to ulcer patients.		
Vand i lungerne Forhøjet blodtryk	9.4	Lægemiddelstof:	Furosemid
		Form:	Tabletter à 40 mg.
		Virkning:	Tvinger væske ud af blodbanen via nyrene til urinblæren, hvorefter væske suges fra lungerne (og andet væv). Virker 1/2 time efter indtagelse. Maksimumeffekt efter 2 timer. Virkningsvarighed: 3 - 6 timer.
		Dosering:	Vand i lungerne: 40 - 120 mg. Kan gentages efter 30 minutter. Blodtryksforhøjelse: 40 - 80 mg x 1 - 3 dagligt. (maksimalt 1500 mg/døgn).
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	5 år fra produktionsdato (Furix®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .

Lung edema High blood pressure	9.4	Active pharmaceutical ingredients:	Furosemide
		Form:	Tablets of 40 mg.
		Effect:	Extracts fluids from the blood vessel via the kidneys to the urinary bladder, after which the fluid is absorbed from the lungs (and other tissue). Takes effect after ½ hour. Maximum effect after 2 hours. Effective for 3-6 hours.
		Dosage:	Lungedema: 40-120 mg. May be repeated after 30 minutes High blood pressure: 40-80 mg once to three times daily (maximum 1500 mg/day).
		Side-effects:	None relevant.
		Validity:	5 years from the date of production (Furix®, Nycomed Denmark).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
Vand i lungerne Forhøjet blodtryk MFAG tabel 2, 9	9.5	Lægemiddelstof:	Furosemid
		Form:	Injektionsvæske 10 mg/ml. Ampuller à 4 ml til intramuskulær eller intravenøs injektion.
		Virkning:	Tvinger væske ud af blodbanen via nyrerne til urinblæren, hvorefter væske suges fra lungerne (og andet væv). Virker inden for 5 minutter efter i.v. injektion. Maksimumeffekt inden 30 minutter.
		Dosering:	Individuelt. 40 - 120 mg. Kan gentages efter 30 minutter.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	5 år fra produktionsdato (Furix®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Lung edema High blood pressure MFAG table 2, 9	9.5	Active pharmaceutical ingredients:	Furosemide
		Form:	Injection liquid 10 mg/ml. Ampoules of 4 ml for intramuscular or intravenous injection.
		Effect:	Extracts fluids from the blood vessel via the kidneys to the urinary bladder, after which the fluid is absorbed from the lungs (and other tissue). Takes effect 5 minutes after i.v. injection. Maximum effect within 30 minutes.
		Dosage:	Dependent on the patient. 40-120 mg. May be repeated after 30 minutes.
		Side-effects:	No relevant ones.
		Validity:	5 years from the date of production (Furix®, Nycomed Denmark).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Forhøjet blodtryk Hjertekrampe Smerter i hjertet Forebyggelse efter blodprop i hjertet	9.6	Lægemiddelstof:	Atenolol
		Form:	Tabletter à 50 mg.
		Virkning:	Sænker puls og blodtryk.
		Dosering:	Forhøjet blodtryk: Voksne. 50-100 mg i døgnet. Hjertekrampe og hjerterytmeforstyrrelser: Sædvanligvis 50 mg 1 gang i døgnet stigende til 100 mg i døgnet. Forebyggelse efter blodprop i hjertet: 50-100 mg 1 gang i døgnet under kontrol af puls og blodtryk.
		Bivirkninger:	Træthed, kvalme, opkastning, diarré, forstoppelse. Langsom puls, sammentrækning af blodkarrene, lavt blodtryk. Muskelsvaghed. Øget svedtendens.
		Holdbarhed:	5 år fra produktionsdato (Atenolol, Orifarm Generics A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes til patienter med astma, kroniske obstruktiv lungesygdom (KOL) og ved svær regulerbar sukkersyge.
High blood pressure Angina pectoris Heart pain Preventiv treatment after Coronary thrombosis	9.6	Active pharmaceutical ingredients:	Atenolol
		Form:	Tablets of 50 mg.
		Effect:	Reduces pulse and blood pressure.
		Dosage:	High blood pressure: Adults: 50- 100 mg daily. Angina pectoris and heart rhythm disturbances: Usually 50 mg once a day, increasing to 100 mg daily. Prevention after coronary thrombosis: 50-100 mg once a day under the control of heart rate and blood pressure.
		Side-effects:	Fatigue, nausea, vomiting, diarrhoea, constipation, low pulse, low blood pressure, contraction of blood vessels, muscle weakness, increased sweating.
		Validity:	5 years from the date of production (Atenolol, Orifarm Generics A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Should not be used in patients with asthma, chronic obstructive pulmonary disease (COPD) and in severe controllable diabetes.

Blodfortyndende	9.8	Lægemiddelstof:	Apixaban
		Form:	Tablet 5 mg
		Virkning:	Virker blodfortyndende. Forhindrer blodpropper.
		Dosering:	5 - 10 mg x 2 dagligt, efter ordination af Radio Medical Danmark
		Bivirkninger:	Øget blødningstendens, kvalme hududslæt, blodig urin, vaginal blødning, blødning under huden.
		Holdbarhed:	3 år fra produktion dato (Blister pk., Teva GmbH) 5 år fra produktions dato (Beholder, Teva GmbH)
		Opbevaring:	Normal temperature
		<i>Anmærkninger:</i>	Søg samråd med Radio Medical.
Blood diluting	9.8	Active Pharmaceutical Ingredients:	Apixaban
		Form:	Tablets 5 mg
		Effect:	Acts as a blood thinner. Prevents blood clots.
		Dosage:	5-10 mg x 2 time daily, after prescription from Radio Medical Denmark
		Side-effects:	Increased tendency to bleed, nausea skin rash, bloody urine, vaginal bleeding, bleeding under the skin.
		Validity:	3 years from production date (Blister pack., Teva GmbH) 5 years from production date (Container, Teva GmbH)
		Storage:	Normale Temperature
		<i>Remarks:</i>	Consult <i>Radio Medical</i>
Stopper blødninger	9.9	Lægemiddelstof:	Tranexamsyre
		Form:	injektion væske 100 mg/ml
		Virkning:	Forebyggelse og behandling af blødninger
		Dosering:	1 g i.v , 3 gange dagligt
		Bivirkninger:	Diarré, kvalme, opkast, hovedpine, allergisk hudreaktion
		Holdbarhed:	3 år fra produktions dato (Tranexamsyre, Pfizer ApS)
		Opbevaring:	Normal temperatur. Må ikke fryses.
		<i>Anmærkninger:</i>	Søg samråd med Radio Medical.
Stops Bleedings	9.9	Active Pharmaceutical Ingredients:	Tranexamacid
		Form:	Injection liquid 100 mg/ml
		Effect:	Prevention and treatment of bleedings
		Dosage:	1g i.v., 3 times daily
		Side-effects:	Diarrhoea, nausea, vomiting, headache, allergic skin reaction
		Validity:	3 years from production date (Tranexamacid, Pfizer ApS)
		Storage:	Normale temperature. Do not freeze
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 10 Hudmidler / Dermatological agents

Svamp Hudsvamp (Se også 14.1)	10.2	Lægemiddelstof:	Miconazol
		Form:	Creme 20 mg/g (2%).
		Virkning:	Dræber svampe i huden.
		Dosering:	Den angrebne hud indsmøres 2 gange dagligt. Behandlingen bør fortsætte 1 uge efter symptomfrihed.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	2 år fra produktionsdato (Brentan®, McNeil Denmark ApS).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Bør ikke anvendes i åbne sår, i hårbund, skæg eller negle.
Fungus Skin fungus (See also 14.1)	10.2	Active pharmaceutical ingredients:	Miconazole
		Form:	Lotion 20 mg/g (2%).
		Effect:	Restricts fungus growth in the skin.
		Dosage:	Apply to the skin twice daily. The treatment should be continued for a week after the symptoms have disappeared.
		Side-effects:	No relevant ones.
		Validity:	2 years from the date of production (Brentan®, McNeil Denmark ApS).
		Storage:	Not above 25°C.
		<i>Remarks:</i>	Should not be used in open wounds, on scalps, beards, or nails.
Lus Fladlus Fnat	10.5	Lægemiddelstof:	Permethrin
		Form:	Creme 50 mg/g (5%).
		Virkning:	Gift, der dræber lus, fnatmidler og deres æg.
		Dosering:	Påsmøres den rene og tørre hud.
		Bivirkninger:	Kløe og rødme af huden.
		Holdbarhed:	5 år fra produktionsdato (Nix®, ACO Hud Nordic AB).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Se instruktionen i indlægsseddel.
Lice Crab lice Scabies	10.5	Active pharmaceutical ingredients:	Permethrin
		Form:	Lotion 50 mg/g (5%).
		Effect:	Poison that kills lice, scab mites, and their eggs.
		Dosage:	Apply to clean, dry skin.
		Side-effects:	Itching and skin blushing.
		Validity:	5 years from the date of production (Nix®, ACO Hud Nordic AB).
		Storage:	Normal temperature.
		<i>Remarks:</i>	See package leaflet.

Grp. 10 Hudmidler / Dermatological agents

(continued)

Lus Fladlus Fnat	10.6	Lægemiddelstof:	Permethrin
		Form:	Shampo 10 mg/g (1%).
		Virkning:	Gift, der dræber lus, fnatmider og deres æg.
		Dosering:	Indgnides i det renvaskede, froterede og fugtige hår og hårbund. Shampooen skal sidde i 10 minutter før den skylles
		Bivirkninger:	Kløe og rødme af huden.
		Holdbarhed:	2 år fra produktionsdato (Nix®, ACO Hud Nordic AB).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Se instruktionen i indlægsseddel.
Lice Crab lice Scabies	10.6	Active pharmaceutical ingredients:	Permethrin
		Form:	Shampoo 10 mg/g (1%).
		Effect:	Poison that kills lice, scab mites, and their eggs.
		Dosage:	Apply to washed, rubbed, and wet hair and scalp. The shampoo should not be rinsed off until after 10 minutes. Dead lice and eggs are removed with a fine comb while the
		Side-effects:	Itching and skin blushing.
		Validity:	2 years from the date of production (Nix®, ACO Hud Nordic AB).
		Storage:	Normal temperature.
		<i>Remarks:</i>	See package leaflet.
Eksem Psoriasis Hudkløe Solskoldning Hudsvi e efter insektb id	10.7	Lægemiddelstof:	Glukokortikoid gruppe II
		Form:	Creme til udvortes brug.
		Virkning:	Modvirker betændelsestilstande, der er fremkaldt af allergiske
		Dosering:	Voksne: Påsmøres tyndt 1-2 gange daglig.
		Bivirkninger:	Hudkløe brændende fornemmelse i huden, tynd hud,
		Holdbarhed:	3 år fra produktionsdato (Locoid, Astellas Pharma A/S).
		Opbevaring:	Almindelig temperatur. Må ikke opbevares koldt eller fryses.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Eczema Psoriasis Itching Sunburns Burning sensation after insect bites	10.7	Active pharmaceutical ingredients:	Glucocorticoid group II
		Form:	Lotion for external use.
		Effect:	Eases inflammations caused by allergic or toxic agents.
		Dosage:	Adults: Apply a thin layer to the skin 1-2 times daily.
		Side-effects:	Pruritic, burning sensation in the skin, thin skin, contact
		Validity:	3 years from the date of production (Locoid, Astellas Pharma A/S).
		Storage:	Normal temperature. Do not refrigerate or freeze.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 10 Hudmidler / Dermatological agents
(continued)

Hudinfektioner	10.8	Lægemiddelstof:	Fusidin
		Form:	Salve 2%
		Virkning:	Bakteriedræbende
		Dosering:	Påsmøres det berørte hudområde 2-3 gange dagligt
		Bivirkninger:	Kontaktseksem
		Holdbarhed:	2 år efter produktion dato (Fusidinsyre, Orifarm)
		Opbevaring:	Normal temperatur
		<i>Anmærkninger:</i>	Søg samråd med Radio Medical Må ikke anvendes i åbne sår.
Skininfections	10.8	Active Pharmaceutical	Fusidin
		Form:	Ointment 2%
		Effect:	Kills bacteria
		Dosage:	Applies the affected skin area 2-3 times daily
		Side-effects:	Contact eczema
		Validity:	2 years from production date (Fusidin acid, Orifarm) 4 weeks after opening.
		Storage:	Normale temperature
		<i>Remarks:</i>	Consult Radio Medical. Do not use in open wounds.

Grp. 11 Desinficerende midler / Disinfectants

Rengøring af: 1. Instrumenter FØR og EFTER brug 2. Hænder 3. Intakt hud	11.1	Lægemiddelstof:	Klorhexidin/ethanol
		Form:	Væske 0,5-5%/70-85%.
		Virkning:	Desinficerende.
		Dosering:	-
		<i>Anmærkninger:</i>	Må ikke anvendes i sår, i øjne eller på slimhinder. Indeholder 85 % alkohol. Se Søfartsstyrelsens Lægebog. Alternativt indkøbes Chlorhexidin 0,5 % og Ethanol minimum 70 %, i 2 separate flasker. Først anvendes chlorhexidin, derefter ethanol.
Disinfection of: 1. Instruments BEFORE and AFTER use 2. Hands 3. Intact skin	11.1	Active pharmaceutical ingredients:	Chlorhexidine/ethanol
		Form:	Liquid 0.5-5%/70-85%.
		Effect:	Disinfective.
		Dosage:	-
		<i>Remarks:</i>	Not be used in wounds, in eyes, or on mucous membranes. Contains 85% alcohol. See the Danish Maritime Authority's Medical Manual. Alternatively, Chlorhexidine 0.5% and ethanol with a minimum of 70% may be purchased in 2 separate bottles. Use chlorhexidine first and then ethanol.
Førstehjælpsbehandling af urene sår Mundskylning	11.2	Lægemiddelstof:	Klorhexidin
		Form:	Væske 0,2%.
		Virkning:	Desinficerende.
		Dosering:	Kan anvendes i forurenede sår, der er utilgængelig for normal sårrensning.
		<i>Anmærkninger:</i>	Se Søfartsstyrelsens Lægebog.
First aid treatment of dirty wounds Mouth rinse	11.2	Active pharmaceutical ingredients:	Chlorhexidine
		Form:	Liquid 0.2%.
		Effect:	Disinfective.
		Dosage:	May be used for dirty wounds that are not accessible for normal wound disinfection.
		<i>Remarks:</i>	See the Danish Maritime Authority's Medical Manual.

Grp. 12 Midler mod mave- og tarmsygdomme / Gastro intestinal agents

Diarré	12.6	Lægemiddelstof:	Loperamid
		Form:	Tabletter à 2 mg.
		Virkning:	Hæmmer tarmenes bevægelser.
		Dosering:	Start: 2 tabletter. Pause 1 time. Derpå 1 tablet for hver uformet afføring. Maksimum: 8 tabletter (16 mg) per døgn.
		Bivirkninger:	Mundtørhed, kvalme, mavesmerter. Risiko for tarmstop.
		Holdbarhed:	5 år fra produktionsdato (Imodium®, McNeil).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Undersøg om den syge har feber. Bør kun anvendes til børn efter <i>Radio Medicals</i> anvisninger. Bør kun anvendes kortvarigt ved akut diarré hos i øvrigt upåvirkede personer.
Diarrhoea	12.6	Active pharmaceutical ingredients:	Loperamide
		Form:	Tablets of 2 mg.
		Effect:	Restricts the motility of the intestines.
		Dosage:	Start: 2 tablets. Wait for 1 hour. 1 tablet for every loose stool. Maximum: 8 tablets (16 mg) daily
		Side-effects:	Dry mouth, nausea, slight stomach pains. Risk of intestinal
		Validity:	5 years from the date of production (Imodium®, McNeil).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Examine whether the patient has a fever. Should only be given to children on directions given by <i>Radio Medical</i> . Use only for a short period of time in case of acute diarrhoea on patients not otherwise affected.
Væsketab Salttab Febertilstande Opkastninger Diarré MFAG tabel 8, 10, 11.	12.7	Lægemiddelstof:	Glucose/kaliumklorid/magnesiumsulfat/natriumcitrat/
		Form:	Afdelte pulvere.
		Virkning:	Erstatter væske og salte.
		Dosering:	1 pulver opløses i 1 liter kogt, afkølet vand. Basisbehov: 2,5 liter per døgn. For hver grad over 37°C gives yderligere 300 ml per døgn. For hver diarré gives yderligere 200 ml per døgn.
		Bivirkninger:	Ingen.
		Holdbarhed:	2 år fra produktionsdato (Revolyt®, Gunnar Kjems APS).
		Opbevaring:	Almindelig temperatur. Færdigblandet opløsning opbevares i køleskab.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .

Grp. 12 Midler mod mave- og tarmsygdomme / Gastro intestinal agents

(continued)

Loss of fluid Loss of salt Feverish conditions Vomiting Diarrhoea MFAG tables 8, 10, 11	12.7	Active pharmaceutical ingredients:	Glucose/potassium/magnesium/sodium citrate/sodium
		Form:	Divided powder.
		Effect:	Replaces liquids and salts.
		Dosage:	1 powder is dissolved in 1 litre of boiled, cooled water. Basic need: 2.5 litres daily. For every degree above 37°C, the patient is given another 300 ml daily. For every instance of diarrhoea, the patient is given another 200 ml daily.
		Side-effects:	None.
		Validity:	2 years from the date of production (Revolyt®, Gunnar Kjems
		Storage:	Normal temperature. The mixed dissolution should be stored in a cold place.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
Mavesår	12.8	Lægemiddelstof:	Omeprazol
		Form:	Tablet eller kapsel 20 à mg.
		Virkning:	Hæmmer dannelsen af mavesyre.
		Dosering:	Mavesår: 20 – 40 mg dagligt. Langtidsbehandling: 10 – 20 mg dagligt.
		Bivirkninger:	Kvalme, opkastning, diarré, forstoppelse, luft i tarmen. Hovedpine, træthed, svimmelhed, søvnforstyrrelser.
		Holdbarhed:	4 år fra produktionsdato (Omeprazol, Copyfarm A/S).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Stomach ulcer	12.8	Active pharmaceutical ingredients:	Omeprazol
		Form:	Tablet or capsules of 20 mg.
		Effect:	Inhibits gastric acid secretion.
		Dosage:	Stomach ulcer: 20-40 mg daily. Long term treatment: 10-20 mg daily.
		Side-effects:	Nausea, vomiting, diarrhoea, constipation, flatulence, headache, fatigue, dizziness, sleep disturbances.
		Validity:	4 years from the date of production (Omeprazol, Copyfarm A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Sure opstød Halsbrand	12.9	Lægemiddelstof:	Antacida
		Form:	Tablet eller oral opløsning.
		Virkning:	Neutraliserer mavesyre.
		Dosering:	Jf. indlægsseddel i præparatet. Efter ordination af <i>Radio Medical</i> .
		Bivirkninger:	Forstoppelse, diarré.
		Holdbarhed:	5 år fra produktionsdato (Alminox®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Acid indigestion Heartburn	12.9	Active pharmaceutical ingredients:	Antacida
		Form:	Tablet or oral solution.
		Effect:	Neutralizes acid in the stomach.
		Dosage:	See patients information leaflet. Prescription from <i>Radio Medical</i> .
		Side-effects:	Constipation, diarrhoea.
		Validity:	5 years from the date of production (Alminox®, Nycomed Danmark).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .
Hæmorider Kløe ved endetarmen Smerter ved afføring	12.10	Lægemiddelstof:	Antihæmorigide creme (skal indeholde glucokortikoid og
		Form:	Salve eller creme.
		Virkning:	Lindring af smerter, hævelse, brændende fornemmelse og kløe i forbindelse med hæmoroider og analrifter.
		Dosering:	Voksne: Smøres på 2-3 gange daglig, samt efter afføring.
		Bivirkninger:	Anal slimhindeatrofi.
		Holdbarhed:	3 år fra produktionsdato (Proctosedyl, 2care4 ApS).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Haemorrhoids Itching at the rectum Pain during defecation	12.10	Active pharmaceutical ingredients:	Anti haemorrhoid ointment (Must contain glucocorticoid and analgesic)
		Form:	Ointment.
		Effect:	Relief of pain, swelling and burning sensation/ itching associated with haemorrhoids, anal lacerations.
		Dosage:	Apply 2-3 times daily and after defecation.
		Side-effects:	Thin skin in the anus.
		Validity:	3 years from the date of production (Proctosedyl, 2care4 ApS).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 12 Midler mod mave- og tarmsygdomme / Gastro intestinal agents

(continued)

Kvalme Opkastninge	12.13	Lægemiddelstof:	Ondansetron
		Form:	Tablet 4 mg
		Virkning:	Modvirker kvalme og opkast
		Dosering:	Efter ordination fra Radio Medical Danmark
		Bivirkninger:	Hovedpine, forstoppelse, Rødme og varmfølelse
		Holdbarhed:	5 år fra produktions dato (Aurobindo Pharma)
		Opbevaring:	Normal temperatur
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i>
Nausea Vomiting	12.13	Active Pharmaceutical Ingredients:	Ondansetron
		Form:	Tablets 4 mg
		Effect:	Prevent nausea and vomiting
		Dosage:	By prescription from Radio Medical Danmark
		Side-effects:	Headache, Constipation, Redness and warm feeling
		Validity:	5 years from production date (Aurobindo Pharma)
		Storage:	Normal temperature
		<i>Remarks:</i>	Consult <i>Radio Medical</i>
Forstoppelse	12.14	Lægemiddelstof:	Movicol®
		Form:	Pulver til oral opløsning
		Virkning:	Binder væske i tarmen, hvorved afføringen blødgøres
		Dosering:	Efter ordination fra Radio Medical Danmark
		Bivirkninger:	Mavesmerter, Anal irritation, Diarré, Flatulens, Kvalme, Meteorisme, Opkastning.
		Holdbarhed:	3 år fra produktions dato (Norgine BV)
		Opbevaring:	Ikke over 25 °C
		<i>Anmærkninger:</i>	Søg samråd med Radio Medical
Constipation		Active Pharmaceutical Ingredients:	Movicol®
		Form:	Powder for oral solution
		Effect:	Binds fluid in the intestine, thereby softening the stools
		Dosage:	By prescription from Radio Medical Danmark
		Side-effects:	Abdominal pain, Anal irritation, Diarrhea, Flatulence, Nausea, Meteorism, Vomiting
		Validity:	3 years from production date (Norgine BV)
		Storage:	Not above 25 °C
		<i>Remarks:</i>	Consult <i>Radio Medical</i>

Grp. 13 Øjen- og øremidler / Agents for Eyes and Ears

Svejsøjne Lokalbedøvelse af øjet MFAG tabel 7	13.2	Lægemiddelstof:	Cinchocain
		Form:	Salve à 5 mg/g.
		Virkning:	Lokalbedøvende.
		Dosering:	1 cm salve bag nederste øjenlåg. Kan gentages efter et par minutter.
		Bivirkninger:	Risiko for maskering af skader eller sygdomme i øjet. Forsinket sårheling.
		Holdbarhed:	2 år fra produktionsdato (Cincain "Optha", Ophta).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.
Arc eye / Welders eye Local anaesthetic of the eye MFAG table 7	13.2	Active pharmaceutical ingredients:	Cinchocaine
		Form:	Ointment of 5 mg/g.
		Effect:	Local anaesthetic.
		Dosage:	Apply 1 cm of ointment behind the lower eyelid. May be repeated after a couple of minutes.
		Side-effects:	Risk of masking injuries or diseases in the eye. Delayed healing of wounds.
		Validity:	2 years from the date of production (Cinchaine, "Optha", Ophta).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.
Allergiske øjensygdomme Inflammatoriske øjensygdomme Øregangseksem	13.3	Lægemiddelstof:	Dexamethason
		Form:	Dråber à 1 mg/ml.
		Virkning:	Standser den allergiske eller inflammatoriske proces.
		Dosering:	Øjet: 1 - 2 dråber bag nederste øjenlåg hver time indtil bedring. Herefter 1 dråbe 4 - 6 gange per døgn. Øret: Øregangen renses, tørres og pakkes med gaze, der er gennemvædet med dexamethason. Gazen holdes fugtig med opløsningen og fjernes efter 24 timer. Behandlingen gentages så længe <i>Radio Medical</i> skønner det nødvendigt.
		Bivirkninger:	Allergi. Risiko for maskering af infektioner, skader eller sygdomme i øjet.
		Holdbarhed:	3 år fra produktionsdato (Maxidex®, S.A. Alcon-couvreur N.V.).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.

Allergic eye diseases Inflammatory eye diseases Ear eczema	13.3	Active pharmaceutical ingredients:	Dexamethasone
		Form:	Eye drops of 1 mg/ml.
		Effect:	Stops the allergic or inflammatory process.
		Dosage:	The eye: 1-2 drops behind the lower eyelid every hour until improvement. Subsequently, 1 drop four-six times daily. The ear: The auditory canal is cleansed and dried, and gauze soaked in dexamethasone is applied. The gauze is kept humid with the dissolution and removed after 24 hours. The treatment is repeated for as long as found necessary by <i>Radio Medical</i> .
		Side-effects:	Allergy. Risk of masking of infections, injuries, or diseases in the eye.
		Validity:	3 years from the date of production produktionsdato (Maxidex®, S.A. Alcon-couvreur N.V.).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.
Grøn stær: Smerter i øjet, Regnbuesyn Tågesyn	13.4	Lægemiddelstof:	Pilocarpin
		Form:	Dråber à 20 mg/ml.
		Virkning:	Nedsætter trykket i det indre øje.
		Dosering:	Hypig drypning bag nederste øjenlåg efter ordination fra <i>Radio Medical</i> .
		Bivirkninger:	Sammentrækning i pupillens lukkemuskel ("knappenålpupil") og hovedpine.
		Holdbarhed:	1½ år fra produktionsdato (Pilocarpin "ophta"®, Ophtha A/S) Må højst anvendes 1 måned efter åbning.
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.
Glaucoma: Pains in the eye, Rainbow vision, Blurred vision	13.4	Active pharmaceutical ingredients:	Pilocarpine
		Form:	Eye drops of 20 mg/ml.
		Effect:	Reduces pressure inside the eye.
		Dosage:	Frequent drops behind the lower eyelid after prescription from <i>Radio Medical</i> .
		Side-effects:	Spasms of the sphincter of the pupil ("pin-hole pupil") and headache.
		Validity:	1½ years from the date of production (Pilocarpin "ophtha"®, Ophtha A/S). To be use within 1 month, from opening the package.
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.

Herpes Helvedesild	13.6	Lægemiddelstof:	Aciclovir
		Form:	Øjensalve à 30 mg/g.
		Virkning:	Hæmmer virus.
		Dosering:	Øjet: 1 cm salve bag nederste øjenlåg 5 gange dagligt indtil 3 døgn efter symptomfrihed. Kan tillige anvendes ved herpesangreb på læberne.
		Bivirkninger:	Øjenirritation
		Holdbarhed:	5 år fra produktionsdato (Zovir®, GlaxoSmithKline Pharma A/ S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Kan ikke anvendes sammen med kontaktlinser.
Herpes Shingles	13.6	Active pharmaceutical ingredients:	Acyclovir
		Form:	Eye ointment of 30 mg/g.
		Effect:	Restricts virus.
		Dosage:	The eye: Apply ointment in a layer of 1 cm behind the lower eyelid five times a day until 3 days after the patient is free of symptoms. May also be used in case of herpes infections of the lips.
		Side-effects:	Irritation of the eye.
		Validity:	5 years from the date of production (Zovir®, GlaxoSmithKline Pharma A/S).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used when wearing contact lenses.
Blødgøring og løsning af ørevoks.	13.9	Lægemiddelstof:	Allantoin
		Form:	Væske.
		Virkning:	Blødgør og løsner ørevoks.
		Dosering:	10-20 dråber dryppes i øret. Et stykke vat anbringes i øret og personen lejres således at dråberne forbliver i øret. Efter 20 -50 min. skylles øret med lunkent vand (37°C).
		Bivirkninger:	Se indlæggessseddel.
		Holdbarhed:	Se produktet(Removax®).
		Opbevaring:	Opbevares ved almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes hvis der er hul i trommehinden. Må ikke anvendes ved betændelse eller smerter i øret og ved allergi overfor indholdsstofferne.
Softens and loosens cerumen	13.9	Active Pharmaceutical ingredients:	Allantoin
		Form:	Fluid.
		Effect:	Softens and dissolving earwax.
		Dosage:	10-20 drops in the ear. A piece of cotton wool is placed in the ear and the person is placed, so that the droplets remain in the ear.
		Side-effects:	See package leaflet.
		Validity:	See expire date on the product (Remowax®).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Do not use if there is a hole in the eardrum. Not for use by inflammation or pain in the ear and allergy to the ingredients.

Inflammatoriske øjensygdomme	13.11	Lægemiddelstof:	Tropicamid
		Form:	Øjendråber 1%.
		Virkning:	Udvider pupillen.
		Dosering:	Akut: 1-2 dråber 3 gange dagligt.
		Bivirkninger:	Feber, forstoppelse, hjertebanken, hallucinationer, excitation,
		Holdbarhed:	3 år fra produktionsdato
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Inflammatory eye diseases	13.11	Active Pharmaceutical ingredients:	Tropicamid
		Form:	Eye drops 1%.
		Effect:	Enlarges the pupil.
		Dosage:	Acut: 1-2 drops 3 times a day.
		Side-effects:	Fever, constipation, palpitations, hallucinations, excitation,
		Validity:	3 years from the date of production
		Storage:	Normal temperatur.
		Remarks:	Consult <i>Radio Medical</i> .
Øregangs eksem	13.12	Lægemiddelstof:	Ciprofloxacin / Hydrocortison
		Form:	Øredråber.
		Virkning:	Behandling af øregangs eksem.
		Dosering:	3 dråber i berørte øre 2 gange daglig.
		Bivirkninger:	Ørekløe, øresmerter, svimmelhed.
		Holdbarhed:	Ubrudt 2 år fra produktionsdato (Ciflox, Novartis Healthcare A/S).
		Opbevaring:	Beskytte mod lys og ikke over 25°C. Må ikke opbevares i
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
Ear eczema	13.12	Active Pharmaceutical ingredients:	Ciprofloxacin / Hydrocortisone
		Form:	Eardrops.
		Effect:	Treatment of ear canal eczema.
		Dosage:	3 drops in affected ear twice daily.
		Side-effects:	Ear itching, earache, dizziness.
		Validity:	Unbroken 2 years from production date (Ciflox, Novartis Healthcare A/S).
		Storage:	Protect from light and not more than 25°C. Do not
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 13 Øjen- og øremidler / Agents for Eyes and Ears

Øjenbetændelse	13.13	Lægemiddelstof:	Kloramfenikol
		Form:	Øjensalve 1%
		Virkning:	Bakteriedræbende
		Dosering:	Smørres 3- 6 gange dagligt i øjet
		Bivirkninger:	Øjenkløe, øjenirritation, allergisk reaktion
		Holdbarhed:	2 år fra produktion dato (Kloramfenikol, 2care4)
		Opbevaring:	2°C til 8°C. Efter anbrud ikke over 25°C
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i>
Eye infection	13.13	Active Pharmaceutical Ingredients:	Chloramphenicol
		Form:	Eye ointment 1%
		Effect:	Kills bacteria
		Dosage:	Apply 3-6 times daily in the eye
		Side-effects:	Eye itching, eye irritation, allergic reaction
		Validity:	2 years from production date (Kloramfenikol, 2care4)
		Storage:	2°C til 8°C. After opening not above 25°C
		<i>Remarks:</i>	Consult <i>Radio Medical</i>

Grp. 14 Gynækologiske midler / Gynaecological agents

Svamp i skeden Hudsvamp (se også 10.2) Neglesvamp	14.1	Lægemiddelstof:	Fluconazol
		Form:	Kapsler à 150 mg.
		Virkning:	Hæmmer svampevækst.
		Dosering:	Skedesvamp: 150 mg som éngangsdosis. Hudsvamp: 150 mg per uge i 4 - 6 uger. Neglesvamp: 150 mg per uge i 3 - 12 måneder (til en ny negl er dannet).
		Bivirkninger:	Allergi. I sjældne tilfælde alvorlig leverforgiftning. Fluconazol påvirker virkningen af adskillige andre medikamenter, herunder bl.a. blodfortyndende og beroligende midler samt p-piller.
		Holdbarhed:	5 år fra produktionsdato (Fluconazol "KRKA", KRKA Sverige AB).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Bør ikke anvendes til gravide.
Vaginal fungus Skin fungus (see also 10.2) Nail fungus	14.1	Active Pharmaceutical ingredients:	Fluconazole
		Form:	Capsule of 150 mg.
		Effect:	Restricts fungal growth.
		Dosage:	Vaginal fungus: 150 mg as a single dose. Skin fungus: 150 mg a week for 4-6 weeks. Nail fungus: 150 mg a week for 3-12 months (until a new nail has been formed).
		Side-effects:	Allergy In rare cases, serious liver poisoning. Fluconazole affects the effect of several other medicaments, including inter alia blood-thinning substances, sedatives, and birth control pills.
		Validity:	5 years from the date of production (Fluconazol "KRKA", KRKA Sverige AB).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . Not to be used by pregnant women.
Blødning EFTER abort eller fødsel	14.2	Lægemiddelstof:	Methylergometrin
		Form:	Tabletter à 0,125 mg.
		Virkning:	Livmodersammentrækkende.
		Dosering:	Efter abort eller fødsel: 0,125 - 0,250 mg 3 gange dagligt.
		Bivirkninger:	Overfølsomhed. Hovedpine, mavesmerter, forhøjet blodtryk, hududslet.
		Holdbarhed:	5 år fra produktionsdato (Methergin®, Novartis).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> . Må ikke anvendes til gravide med blødninger.

Grp. 14 Gynækologiske midler / Gynaecological agents

(continued)

Bleeding AFTER giving birth or miscarriage	14.2	Active Pharmaceutical ingredients:	Methylergometrine
		Form:	Tablets of 0.125 mg.
		Effect:	Contracts the uterus.
		Dosage:	After giving birth or miscarriage: 0.125-0.250 mg three times daily.
		Side-effects:	Allergy, headache, stomachache, high blood pressure, rash.
		Validity:	5 years from the date of production (Methergine®, Novartis).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Not to be used by pregnant women who are bleeding.

Grp. 15. Midler mod forkølelse og astma / Agents for common Cold and Asthma

Hostestillende	15.1	Lægemiddelstof:	Dextromethorphan
		Form:	Tabletter á 30 mg.
		Virkning:	Hostestillende.
		Dosering:	15-30 mg. 3-4 gange daglig.
		Bivirkninger:	Ufrivillige muskelsammentrækning, dødsighed, svimmelhed, hallucinationer, mani.
		Holdbarhed:	3 år fra produktionsdato (Dexofan® Nycomed).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Bør ikke anvendes af børn under 6 år. Bør ikke anvendes ved samtidig behandling med 7.6 og 7.7 Metronidazol. Bør ikke anvendes ved samtidig behandling med Disulfiram (Antabus®).
Cough	15.1	Active Pharmaceutical ingredients:	Dextromethorphan
		Form:	Tablets of 30 mg.
		Effect:	Reduces coughs.
		Dosage:	15 – 30 mg. 3-4 times daily.
		Side-effects:	Unvoluntary muscle contraction, drowsiness, dizziness, hallucinations and mania.
		Validity:	3 years from the date of production (Dexofan®, Nycomed).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . Should not be used by children under the age of 6 years. Should not be used during concomitant treatment with 7.6 or 7.7 Metronidazole. Should not be used during concomitant treatment with disulfiram (Antabuse®).

Grp. 15 Midler mod forkølelse og astma / Agents for common Cold and Asthma (continued)

Forkølelse Bihulebetændelse Kæbehulebetændelse Pandehulebetændelse Mellemørebetændelse	15.5	Lægemiddelstof:	Xylometazolin
		Form:	Næsedråber/næsespray. 1 mg/ml.
		Virkning:	Skaffer afløb fra mellemøret, bihuler, pandehuler og kæbehuler. Virker øjeblikkeligt og i 10 - 12 timer.
		Dosering:	2 pust/dråber i hvert næsebor 2 - 3 gange dagligt i maksimum
		Bivirkninger:	Ved længere tids anvendelse kan stoffet skade næsens
		Holdbarhed:	3 år fra produktionsdato (Zymelin®, Nycomed Danmark).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Må ikke anvendes ved symptomer på grøn stær (se 13.4).
Cold Sinusitis Maxillary sinusitis Frontal sinusitis Otitis media	15.5	Active Pharmaceutical ingredients:	Xylometazoline
		Form:	Nose drops/nose spray. 1 mg/ml.
		Effect:	Opens drainage from middle ears and sinuses. Takes effect immediately and stays effective for 10-12 hours.
		Dosage:	2 sprays/drops in each nostril two-three times daily for a
		Side-effects:	If used for a long period, the substance may damage the
		Validity:	3 years from the date of production (Zymelin®, Nycomed Danmark).
		Storage:	Normal temperature.
		Remarks:	Not to be used if there are symptoms of glaucoma (see 13.4).
Førstevalg ved astma eller astmalignende vejtrækning som følge af udsættelse for røg, gasarter og dampe.	15.7	Lægemiddelstof:	Salbutamol.
		Form:	Spray, 0.1 mg per dose.
		Virkning:	Løser kramper i den glatte muskulatur i luftvejene.
		Dosering:	1 -2 dosage as required. Maximum 4 times daily
		Bivirkninger:	Muskelkramper, hurtig puls, rystelser, hovedpine
		Holdbarhed:	3 years from the date of production (Salbutamol, Teva B.V.)
		Opbevaring:	Ikke over 25 °C - beskyt mod kulde og frost
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> .
First choice of medicament in case of asthma or asthma-like breathing caused by exposure to smoke, gases, and fumes.		Active Pharmaceutical Ingredients:	Salbutamol
		Form:	Spray, 0.1 mg per dose.
		Effect:	Loosens convulsions in the smooth musculature in the
		Dosage:	1 -2 dosage as required. Maximum 4 times daily
		Side-effects:	Muscle cramps, fast pulse, tremors, headache
		Validity:	3 years from the date of production (Salbutamol, Teva B.V.)
		Storage:	Below 25 °C - protect against cold and frost
		Remarks:	Consult <i>Radio Medical</i> .

Grp. 15 Midler mod forkølelse og astma / Agents for common Cold and Asthma (continued)

Astma eller astmalignende vejrtrækning som følge af udsættelse for røg, gasarter og dampe. IMFAG tabel 9	15.8	Lægemiddelstof:	Glucocorticoid til inhalation
		Form:	Spray 100-200 mikrogram/dosis
		Virkning:	Fjerner inflammatorisk ødem og hævelse i luftvejene.
		Dosering:	1 dosis efter behov.
		Bivirkninger:	Svamp i munden, hovedpine, hæshed, hoste og halsgener
		Holdbarhed:	3 år fra produktionsdato (AeroBec, 2care4)
		Opbevaring:	Ikke over 25 °C
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Asthma or asthma-like breathing caused by exposure to smoke, gases, and fumes. MFAG tabel 9	15.8	Active Pharmaceutical Ingredients:	Glycocorticoid for inhalation
		Form:	Spray 100-200 microgram/dosage
		Effect:	Removes inflammatory oedema and swelling of the respiratory tract.
		Dosage:	1 dose as required
		Side-effects:	Fungus in the mouth, headache, hoarseness, cough and sore throat
		Validity:	3 years from the date of production (AeroBec, 2care4)
		Storage:	Below 25 °C
<i>Remarks:</i>	Consult <i>Radio Medical</i> .		

Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar			
1. valgs præparat ved: Blodtab Væsketab Forebyggelse af shock MFAG tabel 7	16.1	Lægemiddelstof:	Natriumklorid
		Form:	Infusionsvæske 9 mg/ml. Beholdere à 1000 ml til intravenøs anvendelse.
		Virkning:	Erstatter væske og salte.
		Dosering:	Normalt ikke over 500 ml per time (godt 150 dråber per minut). 20 dråber = 1 ml.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	3 år fra produktionsdato (Natriumklorid isotonisk, Frenius Kabi).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved malaria og ved hjertetilfælde bør infusionshastigheden være langsom af hensyn til risikoen for udvikling af lungeødem (9.4/9.5).
First choice of medicament in case of: Loss of blood Loss of liquid Prevention of shock MFAG table 7	16.1	Active Pharmaceutical ingredients:	Sodium chloride
		Form:	Infusion liquid 9 mg/ml. Containers of 1000 ml for intravenous use.
		Effect:	Replaces liquids and salts.
		Dosage:	Normally, no more than 500 ml per hour (about 150 drops a minute). 20 drops = 1 ml.
		Side-effects:	None relevant.
		Validity:	3 years from the date of production (Sodium chloride, Frenius Kabi).
		Storage:	Normal temperature.
		Remarks:	Consult <i>Radio Medical</i> . In case of malaria and heart attacks, the infusion speed should be slow because of the risk of developing lung oedema (9.4/9.5).
Ernæringserstatning Væsketab (Se også 12.7)	16.2	Lægemiddelstof:	Glucose og natriumklorid
		Form:	Infusionsvæske med glucose 25 mg/ml og natriumklorid 4,5 mg/ml. Beholdere à 1000 ml til intravenøs anvendelse.
		Virkning:	Erstatter væske, glucose og salte.
		Dosering:	Normalt ikke over 500 ml per time (godt 150 dråber per minut). 20 dråber = 1 ml.
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	3 år fra produktionsdato (Natriumklorid-Glucose "Braun", B. Braun Medical A/S).
		Opbevaring:	Almindelig temperatur.
		Anmærkninger:	Søg samråd med <i>Radio Medical</i> . Ved malaria og ved hjertetilfælde bør infusionshastigheden være langsom af hensyn til risikoen for udvikling af lungeødem (9.4/9.5).

Food substitute Loss of liquids (See also 12.7)	16.2	Active Pharmaceutical ingredients:	Glucose and sodium chloride
		Form:	Infusion liquid containing glucose 25 mg/ml and sodium chloride 4.5 mg/ml. Containers of 1000 ml for intravenous use.
		Effect:	Replaces liquids, glucose and salts.
		Dosage:	Normally, no more than 500 ml per hour (about 150 drops a minute). 20 drops = 1 ml.
		Side-effects:	None relevant.
		Validity:	3 years from the date of production (Natriumklorid-Glucose" Braun", B. Braun Medical A/S).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> . In case of malaria and heart attacks, the infusion speed should be slow because of the risk of developing lung oedema (9.4/9.5).
Væske til gennemskylning af intravenøs kanyle. Sårbehandling. Øjenbehandling	16.5	Lægemiddelstof:	Natriumklorid
		Form:	Væske 0,9% i ca. 20 ml beholder.
		Virkning:	Fjerner blod fra venekanyler, rensesår og øjne.
		Dosering:	Se instruks i Søfartsstyrelsens lægebog eller efter ordination fra <i>Radio Medical</i> .
		Bivirkninger:	Ingen relevante.
		Holdbarhed:	3 år fra produktionsdato (Fresenius Kabi).
		Opbevaring:	Almindelig temperatur.
		<i>Anmærkninger:</i>	Søg samråd med <i>Radio Medical</i> .
Fluid for flushing intravenous needle. Treatment of wounds. Treatment of eyes	16.5	Active Pharmaceutical ingredients:	Sodium chloride
		Form:	Fluid 0,9 % in containers of 20 ml.
		Effect:	Removes blood from intravenous needle, clean wounds and eyes.
		Dosage:	See the instructions in The Maritime Medical Manual or on prescription from <i>Radio Medical</i> .
		Side-effects:	No relevant ones.
		Validity:	3 years from the date of production (Fresenius Kabi).
		Storage:	Normal temperature.
		<i>Remarks:</i>	Consult <i>Radio Medical</i> .

Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar

(continued)

Vand til opløsning af 7.9 Ceftriaxon pulver	16.7	Lægemiddelstof:	Sterilt vand
		Form:	Væske i 10 ml beholder
		Virkning:	-
		Dosering:	-
		Bivirkninger:	-
		Holdbarhed:	3 år fra produktionsdato (Sterilt vand Mini-plasco®, B. Braun Melsungen AG)
		Opbevaring:	Normal temperatur
		<i>Anmærkninger:</i>	-
Water for dissolution of 7.9 Ceftriaxon powder		Active Pharmaceutical Ingredients:	Sterile water
		Form:	Liquid in 10 ml container
		Effect:	-
		Dosage:	-
		Side-effects:	-
		Validity:	3 years from the date of production (sterile water mini-plasco®, B. Braun Melsungen AG)
		Storage:	Normale temperature
		<i>Remarks:</i>	-

**Fortegnelse og kontroldokument over
*lægemiddelstoffer***

**Inventory and Control Document
*of Active Pharmaceutical Ingredients***

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A2
Grp. 1 Ilt / Oxygen						
1.1	V03AN01	Medicinsk ilt	Gas	-	20 l / 200 bar	20 l / 200 bar
1.1	V03AN01	Oxygen, Medical	Gas	-	20 l / 200 bar	20 l / 200 bar
1.3	V03AN01	Medicinsk ilt passende til A.1 (se note)	Gas	-	2 x 2 l / 200 bar	0
1.3	V03AN01	Oxygen Medical suitable to A.1 (see note)	Gas	-	2 x 2 l / 200 bar	0
1.3 Meddelelser A kapitel A IX B, bilag 1 / Notice A Chapter IX B, annex 1						
Grp. 2 Lokalbedøvelse / Local Anaesthetic						
2.1(π)	N01BB02	Lidocain	Inj. Væske	10 mg / ml	5 x 20 ml	0
2.1(π)	N01BB02	Lidocain	Inj. Liquid	10 mg / ml	5 x 20 ml	0
2.2	D04AB01	Lidocain	Gel	2%	2 x 30 g	0
2.2	D04AB01	Lidocain	Gel	2%	2 x 30 g	0
Grp. 3 Smertestillende midler / Analgesics						
3.1	N02BE01	Paracetamol	Tablet	500 mg	200 stk.	0
3.1	N02BE01	Paracetamol	Tablets	500 mg	200 pcs.	0
3.2(π)	M01AE01	Ibuprofen	Tablet	400 mg	100 stk.	0
3.2(π)	M01AE01	Ibuprofen	Tablets	400 mg	100 pcs.	0
3.4 (π)	N02AA01	Morfin Δ	Inj. Væske	10 mg / ml	10 x 1 ml	0
3.4 (π)	N02AA01	Morphine Δ	Inj. Liquid	10 mg / ml	10 x 1 ml	0
3.7	M02AA15	Diclofenac	Gel	11,6 mg / g	50 g	0
3.7	M02AA15	Diclofenac	Gel	11,6 mg / g	50 g	0
3.8 (π)	M01AB05	Diclofenac	Supp.	50 mg	50 stk.	0
3.8 (π)	M01AB05	Diclofenac	Supp.	50 mg	50 pcs.	0
3.9(π)	N02AA01	Morphin Δ	Tablet	10 mg	30 stk	0
3.9(π)	N02AA01	Morphine Δ	Tablets	10 mg	30 pcs.	0
Grp. 4 Allergi, shock og søsyge / Allergy, Shock and Seasickness						
4.1(π)	C01CA24	Adrenalin	Inj. Væske	1 mg / ml	5 x 1ml	0
4.1(π)	C01CA24	Epinephrine	Inj. Liquid	1 mg / ml	5 x 1ml	0
4.2	R06AX18	Acrivastine	Kapsel	8 mg	12 pcs.	0
4.2	R06AX18	Acrivastine	Capsula	8 mg	12 pcs.	0
4.3(π)	H02AB09	Hydrocortison	Inj. Tørstof + væske	100 mg	2 hætteglas	0
4.3(π)	H02AB09	Hydrocortisonsuccinate	Inj. Subst + solvens	100 mg	2 vials	0
4.5(π)	N07CA02	Cinnarizine Δ	Tablet	25 mg	50 pcs.	0
4.5(π)	N07CA02	Cinnarizine Δ	Tablets	25 mg	50 pcs.	0
4.6(π)	H02AB06	Prednisolon	Tablet	25 mg	30 stk.	0
4.6(π)	H02AB06	Prednisolon	Tablets.	25 mg	30 pcs.	0

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A2
Grp. 5 Antidoter / Antidotes						
5.1(π)	V03AB15	Naloxon	Inj. Væske	0,4 mg /ml	5 x 1 ml	0
5.1(π)	V03AB15	Naloxon	Inj. Liquid	0,4 mg /ml	5 x 1 ml	0
5.2(π)	H04AA01	Glucagon "K" (se note)	Inj. Tørstof + væske	1 mg	0	0
5.2(π)	H04AA01	Glucagon "R" (see note)	Inj. Subst + solvens	1 mg	0	0
5.3(π)	A03BA01	Atropine	Inj. Væske	1mg/ml	0	15 x 1ml
5.3(π)	A03BA01	Atropine	Inj. Liquid	1mg/ml	0	15 x 1ml
5.4(π)	D11AX03	Calcium gluconate	Gel	2%	0	5 x 25 g
5.4(π)	D11AX03	Calcium gluconate	Gel	2%	0	5 x 25 g
5.5(π)	A12AA02 / A12AA06	Calcium lactogluconate	Brusetablet	500 mg	0	40 stk.
5.5(π)	A12AA02 / A12AA06	Calcium lactogluconate	Effervescent tablet	500 mg	0	40 pcs.
5.6	A07BA01	Aktivt kul	Pulver	50 g	0	2 x 50 g
5.6	A07BA01	Charcoal, activated	Powder	50 g	0	2 x 50 g
5.7	-	Ethyl alkohol	Væske	99.9%	0	500 ml
5.7	-	Ethyl alcohol	Liquid	99.9%	0	500 ml
5.2: 2 sæt skal medbringes af besætningsmedlemmer med insulinkrævende sukkersyge / 2 sets to be carried by crew members with insulin dependent diabetes.						
Grp. 6 Akut sindssyge og kramper / Mental disorder and convulsion						
6.2(π)	N05BA01	Diazepam Δ	Tablet	5 mg	25 stk.	0
6.2(π)	N05BA01	Diazepam Δ	Tablets	5 mg	25 pcs.	0
6.3(π)	N05BA01	Diazepam Δ	Klysma	4 mg / ml	10 x 2½ ml	0
6.3(π)	N05BA01	Diazepam Δ	Enema	4 mg / ml	10 x 2½ ml	0
6.4(π)	N05AD01	Haloperidol Δ	Inj. Væske	5 mg / ml	10 x 1 ml	0
6.4(π)	N05AD01	Haloperidol Δ	Inj. Liquid	5 mg / ml	10 x 1 ml	0
6.5(π)	N05CF02	Zolpidem Δ	Tablet	10 mg	20 stk.	0
6.5(π)	N05CF02	Zolpidem Δ	Tablets	10 mg	20 pcs.	0

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A2
Grp. 7 Infektioner / Infections						
7.1(π)	J01CE02	Phenoxymethylpenicillin	Tablet	660-800 mg	100 stk.	0
7.1(π)	J01CE02	Phenoxymethylpenicillin	Tablets	660-800 mg	100 pcs.	0
7.2(π)	J01CA04	Amoxicillin	Tablet	500 mg	30 stk.	0
7.2(π)	J01CA04	Amoxicillin	Tablets	500 mg	30 pcs.	0
7.6(π)	P01AB01	Metronidazol	Tablet	500 mg	24 pcs.	0
7.6(π)	P01AB01	Metronidazole	Tablets	500 mg	24 pcs.	0
7.2(π)	J01CA04	Amoxicillin	Tablet	500 mg	30 stk.	0
7.2(π)	J01CA04	Amoxicillin	Tablets	500 mg	30 pcs.	0
7.7(π)	P01AB01	Metronidazol	Supp.	500 mg	20 stk.	0
7.7(π)	P01AB01	Metronidazole	Supp.	500 mg	20 pcs.	0
7.8(π)	J01MA02	Ciprofloxacin	Tablet	250 mg	40 stk.	0
7.8(π)	J01MA02	Ciprofloxacin	Tablets	250 mg	40 pcs.	0
7.9(π)	J01DD04	Ceftriaxon (se note)	inj. Tørstof	1 g	20 x 1g	0
7.9(π)	J01DD04	Ceftriaxone (see note)	inj. Subst	1 g	20 x 1g	0
7.10 (π)	-	Tetanus holdig vaccine "K"	Inj. Væske	-	1 stk.	0
7.10 (π)	-	Vaccine containing Tetanus "R"	Inj. Liquid	-	1 pc.	0
7.11(π)	J06BB02	Tetanus immunglobulin "K"	Inj. Væske	250 IU.	1 x 1 ml	0
7.11(π)	J06BB02	Tetanus immunglobulin "R"	Inj. Liquid	250 IU.	1 x 1 ml	0
7.12(π)	J01CF05	Flucloxacillin	Tablet	500 mg	50 pcs	0
7.12(π)	J01CF05	Flucloxacillin	Tablet	500 mg	50 pcs	0
7.15(π)	J01FA10	Azithromycin	Tablet	500 mg	5 x 3 stk.	0
7.15(π)	J01FA10	Azithromycin	Tablets	500 mg	5 x 3 pcs.	0
7.16(π)	J01CA08	Pivmecillinam	Tablet	400 mg	50 stk.	0
7.16(π)	J01CA08	Pivmecillinam	Tablets	400 mg	50 pcs.	0
7.17(π)	J05AB01	Aciclovir	Tablet	400 mg	70 stk.	0
7.17(π)	J05AB01	Acyclovir	Tablets	400 mg	70 pcs.	0

ADVARSEL: I DK anvendes Ceftriaxon (7,9) i stedet for den i MFAG angivet Cefuroxim.

WARNING: In DK is Ceftriaxone (7.9) used instead of Cefuroxime as described in the MFAG.

7.9 Skibe med en overfartstid under 24 timer kan reducere mængden af Ceftriaxon fra 20 x1g til 5 x1 g.

7.9 Ships which are in engaged in voyage less than 24 hours may reduce the amount of Ceftriaxon from 20 x1g to 5 x1 g

Grp. 8 Malaria / Malaria (se note /see note)					Treatment	Prevention
8.1(π)	P01BB51	Atovaquon/Proguanil	Tablet	250 / 100 mg	0	24 stk./ besætningmedl
8.1(π)	P01BB51	Atovaquon/Proguanil	Tablet	250 / 100 mg	0	24 pcs./ crewmember
8.2(π)	J01AA02	Doxycyclin	Tablet	100 mg	0	80 stk.
8.2(π)	J01AA02	Doxycycline	Tablet	100 mg	0	80 pcs.
8.4(π)	P01BF01	Artemether og lumefantrine	tbl.	20 mg / 120 mg	24 stk	0
8.4(π)	P01BF01	Artemether and lumefantrine	tablets.	20 mg / 120 mg	24 pcs	0

Malaria midler til behandling skal forefindes i skibe, der sejler til områder med malaria - uanset risiko.

Medicinsk forebyggelse kræves hovedsagelig ved sejlads i høj risikoområder.

Højrisikoområder findes opdateret og beskrevet i Søfartsstyrelsens malariastrategi på følgende link

<http://www.soefartsstyrelsen.dk>.

Drugs for treatment of malaria must be carried by all ships sailing to or in areas where malaria is recorded - irrespective of the risk.

Medical prevention is mainly required for ships sailing in high risk areas.

The High risk areas are available and updated on the "Danish Maritime Authority's malaria strategy" on the following link

<http://www.dma.dk>

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A2
Grp. 9 Hjerteog kredsløb / Cardiac and circuity						
9.1	C01DA02	Glycerylnitrat	Mund pray	0,4 mg/dose	1x200 doser	0
9.1	C01DA02	Glyceryl trinitrate	Oral spray	0,4 mg/dose	1x200 doses	0
9.3	B01AC06	Acetylsalisylsyre	Tablet	150 mg	100 stk.	0
9.3	B01AC06	Acetylsalicylic Acid	Tablet	150 mg	100 pcs.	0
9.4(π)	C03CA01	Furosemid	Tablet	40 mg	50 stk.	0
9.4(π)	C03CA01	Furosemide	Tablet	40 mg	50 pcs.	0
9.5(π)	C03CA01	Furosemid	Inj. væske	10 mg/ml	10x4 ml	0
9.5(π)	C03CA01	Furosemide	Inj. Liquid	10 mg/ml	10x4 ml	0
9.6(π)	C07AB03	Atenolol	Tablet	50 mg	100 stk.	0
9.6(π)	C07AB03	Atenolol	Tablet	50 mg	100 pcs.	0
9.8(π)	B01AF02	Apixaban	Tablet	5 mg	28 stk.	0
9.8(π)	B01AF02	Apixaban	Tablets	5 mg	28 pcs.	0
9.9(π)	B02AA02	Tranexamsyre	Inj. væske	100 mg/ml	10 x 5 ml	0
9.9(π)	B02AA02	Tranexamsyre	Inj. liquid	100 mg/ml	10 x 5ml	0
Grp. 10 Hud / Skin						
10.2	D01AC02	Miconazol	Creme	2%	4 x15 g	0
10.2	D01AC02	Miconazol	Lotion/Cream	2%	4 x15 g	0
10.5	P03AC04	Permethrin	Creme	5%	30 g	0
10.5	P03AC04	Permethrin	Lotion/Cream	5%	30 g	0
10.6	P03AC04	Permethrin	Shampoo	1%	50 ml	0
10.6	P03AC04	Permethrin	Shampoo	1%	50 ml	0
10.7	D07AB02	Glucocorticoid gruppe II	Creme	Gruppe II	2 x 30 g	0
10.7	D07AB02	Glucocorticoid group II	Lotion/Cream	Group II	2 x 30 g	0
10.8(π)	D06AX01	Fusidin	Cream / Salve	2%	4 x15 g	0
10.8(π)	D06AX01	Fusidine	Cream / Lotion	2%	4 x 15 g	0
Grp. 11 Desinficering / Desinfectants						
11.1	-	Klorhexidin/ethanol (se note)	Væske	0.5- 5%/70-85%	175 ml	0
11.1	-	Chlorhexidine/ethanol (see note)	Liquid	0.5-5%/70-85%	175 ml	0
11.2	-	Klorhexidin	Væske	0.2 %	4 x 250 ml	0
11.2	-	Chlorhexidine	Liquid	0.2 %	4 x 250 ml	0
11.1: Alternativt kan anvendes Klorhexidin 0,5-5% og Ethanol minimum 70%, i 2 separate flasker. Blandes 1:1 Alternatively, Chlorhexidine 0,5-5% and ethanol with a minimum of 70% in 2 separate bottles. Mix 1:1						

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A2
Grp. 12 Mave og tarm / Gastrointestinal						
12.6	A07DA03	Loperamid	Tablet	2 mg	60 pcs.	0
12.6	A07DA03	Loperamide	Tablet	2 mg	60 pcs.	0
12.7	-	Glukose/kalium/ natriumcitrat /natriumklorid	Pulver	13,5/1,5/2,9/ 2,6 g/pulver	18 x 20,5 g	0
12.7	-	Glucose/potassium/sodiumcitrate /sodiumchlorid	Powder	13,5/1,5/2,9/ 2,6 g/powder	18 x 20,5 g	0
12.8(π)	A02BC01	Omeprazol	Tablet/Kapsel	20 mg	28 pcs.	0
12.8(π)	A02BC01	Omeprazol	Tablet/Capsula	20 mg	28 pcs.	0
12.9	A02A	Antacida	Tablet eller oral opløsning	-	40 stk. eller 250 ml	0
12.9	A02A	Antacid	Tablet or oral solution	-	40 pcs. or 250 ml	0
12.10	C05A	Hæmoride creme (skal indeholde glukokortikoid and smertestillende)	Creme/Salve	-	2 x 30 g	0
12.10	C05A	Anti hæmorrhoid agent (Must contain glucocorticoid and analgesi)	Lotion/Ointment	-	2 x 30 g	0
12.13	A04AA01	Ondasetron	Tablet	4 mg	10 stk.	0
12.13	A04AA01	Ondasetron	tablets.	4 mg	10 pcs.	0
12.14	A06AD65	Movicol	oral opløsning	46,6 mg kaliumchlorid, 13,125 g macrogol 3350, 350,7 mg natriumchlorid , 178,5 mg natriumhydro gencarbonat.	4 bokse af 8 stk.	0
12.14	A06AD65	Movicol	oral solution	46,6 mg kaliumchlorid, 13,125 g macrogol 3350, 350,7 mg natriumchlorid , 178,5 mg natriumhydro gencarbonat.	4 boxes of 8 pcs.	0
12.13 Erstatte, den i MFAG nævnte, injektion Metroclorpramid / 12.13 Replaces the injection Metroclorpramide described in MFAG.						

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Styrke / Strength	Mængde/ Quantity A	MFAG A2
Grp. 13 Øjne og øre / Eyes and ears						
13.2(π)	S01HA06	Cinchocain	Salve	0,50 %	5 x 1g	0
13.2(π)	S01HA06	Cinchocaine	Ointment	0,50 %	5 x 1g	0
13.3(π)	S01BA01	Dexamethason	Dråber	1 mg / ml	5 ml	0
13.3(π)	S01BA01	Dexamethasone	Drops	1 mg / ml	5 ml	0
13.4(π)	S01EB51	Pilocarpin	Dråber	2 %	10 ml	0
13.4(π)	S01EB51	Pilocarpine	Drops	2 %	10 ml	0
13.6(π)	S01AD03	Aciclovir	Salve	3 %	4,5 g	0
13.6(π)	S01AD03	Acyclovir	Ointment	3 %	4,5 g	0
13.9	-	Medium til at blødgøre ørevoks, inkl. Øresprøjte	-	-	10 ml	0
13.9	-	Medium to soften earwax, incl. Earsyringe	-	-	10 ml	0
13.11(π)	S01FA06	Tropicamid	Dråber	1 %	15 ml	0
13.11(π)	S01FA06	Tropicamide	Drops	1 %	15 ml	0
13.12(π)	S02CA03	Ciprofloxacin / Hydrocortison	Øredråber	10+2 mg/ml	3 x 10 ml	0
13.12(π)	S02CA03	Ciprofloxacin / Hydrocortisone	Eardrops	10+2 mg/ml	3 x 10 ml	0
13.13(π)	S01AA01	Kloramfenikol	øjensalve	1%	2 x 4,3 - 5 g	0
13.13(π)	S01AA01	Chloramphenicol	eye ointment	1%	2 x 4,3 - 5 g	0
Grp. 14 Gynækologi / Gynaecology						
14.1(π)	J02AC01	Fluconazol	Kapsel	150 mg	2 stk.	0
14.1(π)	J02AC01	Fluconazole	Capsula	150 mg	2 pcs.	0
14.2(π)	G02AB01	Methylergometrin	Tablet	0,125 mg	10 pcs.	0
14.2(π)	G02AB01	Methylergometrine	Tablet	0,125 mg	10 pcs.	0
14.2 skal kun forefindes, HVIS der er kvinder om bord. 14.2 shall only be available IF women on board.						
Grp. 15 Astma og forkølelse / Astma and common cold						
15.1	R05DA09	Dextromethorphan	Tablet	30 mg	40 stk.	0
15.1	R05DA09	Dextromethorphane	Tablet	30 mg	40 pcs.	0
15.5	R01AA07	Xylometazolin	Dråber / Spray	1 mg / ml	3 x 10 ml	0
15.5	R01AA07	Xylometazoline	Drops / Spray	1 mg / ml	3 x 10 ml	0
15.7(π)	R03AC03	Salbutamol	Spray	0,1 mg/dosis	200 doser	0
15.7(π)	R03AC03	Salbutamol	Spray	0,1 mg/dosage	200 dosis	0
15.8(π)	R03BA01	Glucocorticoid (inhalation)	Spray	100-200 mikrogram pr. dosis	100 doser	4 x 100 doser
15.8(π)	R03BA01	Glucocorticoid (inhalation)	Spray	100-200 mikrogram per dosage	100 dosis	4 x 100 dosis
Grp. 16 Infusionsvæsker og lign. / Liquid for infusion and similar (se note / see note)						
16.1(π)	B05BB01	Natriumklorid	Inf. Væske	0.9 %	10 x 1 liter	0
16.1(π)	B05BB01	Sodium chloride	Inf. Liquid	0.9 %	10 x 1 litre	0
16.2(π)	B05BB02	Glucose + Natriumklorid	Inf. Væske	25/4,5 mg/ml	10 x 1 liter	0
16.2(π)	B05BB02	Glucose + Sodium chlorid	Inf. Liquid	25/4,5 mg/ml	10 x 1 litre	0
16.5	V07AB	Natriumklorid 20 ml	Væske	0.9 %	20 x 20 ml	0
16.5	V07AB	Sodium chloride 20 ml	Liquid	0.9 %	20 x 20 ml	0
16.7	V07AB	Steril vand til blanding af 7.9	Parenteral væske	-	20 x 10 ml	0
16.7	V07AB	Sterile water to mix 7.9	Parenteral væske	-	20 x 10 ml	0
Infusionsvæskerne må ikke være i glasbeholdere Infusion liquids may not be in glass container						

Besætningens medbragte medicin / Crew member's private medicine

Tilhører besætningsmedlem <i>Belongs to crewmember</i>	Medicinens navn <i>Name of medicine</i>	Antal <i>Quantity</i>	Daglig dosis <i>Daily dosage</i>	Dato ind Date <i>in</i>	Kvittering modtagelsen <i>Signature receiving</i>	Dato ud <i>Date out</i>	Kvittering for udlevering <i>Signature for extradition</i>

Besætningens medbragte medicin / Crew member's private medicine

Tilhører besætningsmedlem Belongs to crewmember	Medicinens navn Name of medicine	Antal Quantity	Daglig dosis Daily dosage	Dato ind Date in	Kvittering modtagelsen Signature recieving	Dato ud Date out	Kvittering for udlevering Signature for extradition

**Fortegnelse og Kontrolldokument
over medicinsk udstyr**

**Inventory and Control Document
of Medical Equipment**

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A2
A. Genoplivningsudstyr / Resuscitation equipment			
A.1	Mobilt genoplivningsudstyr, klargjort (se note)	1 sæt	0
A.1	Mobile resuscitation equipment (see note)	1 set	0
A.4	Tungeholdere, størrelse 2	0	2 stk.
A.4	Guidel airway, size 2	0	2 pcs.
A.5	Tungeholdere, størrelse 3	1 stk.	1 stk.
A.5	Guidel airway, size 3	1 pcs.	1 pcs.
A.6	Tungeholdere, størrelse 4	0	2 stk.
A.6	Guidel airway, size 4	0	2 pcs.
A.7	Ekstra ventilationspose med ventil og voksenmaske	0	1 stk.
A.7	Extra ventilation bag with valve and mask	0	1 pcs.
A.8	Kraftig klædesaks	1 stk.	0
A.8	Strong scissors for clothes	1 pcs.	0
A.9	Sugekatetre, lige, CH 12	5 stk.	0
A.9	Suction catheters, straight, CH 12	5 pcs.	0
A.10	Tæppe til behandling af hypotermi	1 stk.	0
A.10	Blanket for treatment of hypothermia	1 pcs.	0
A.11	Almindelige tæpper	6 stk.	0
A.11	Ordinary blankets	6 pcs.	0
A.12	Saturationsmåler (fingermodel)	1 stk.	0
A.12	Puls oximeter (finger)	1 pcs.	0
A.13	Hudson maske med reservoirpose med ca. 1,8 m slange	5 stk.	5 stk.
A.13	Hudson mask with reservoir bag with approx. 1.8 m tube	5 pcs.	5 pcs.
A.18	Hjertestarter (AED)	1 stk.	0
A.18	Automatic extern defibrillator (AED)	1 pc.	0
A.19	Iltbrille	5 stk.	0
A.19	Nasal Cannula	5 pcs.	0
A.1: Meddelelser A kapitel A IX B, bilag 1 / A.1: Notice A Chapter IX B, annex 1			

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A2
B. Forbindinger og plastre / Dressings and adhesive tape			
B.1	Sterile gazekompresser, ca.5 x 5 cm, ca.8 lag	20 stk.	0
B.1	Sterile gaze compress, approx.5 x 5 cm, approx. 8 layers	20 pcs.	0
B.2	Sterile gazekompresser, ca. 10 x 10 cm, ca. 8 lag	20 stk.	0
B.2	Sterile gaze compress, approx.10 x10 cm, approx. 8 layers	20 pcs.	0
B.3	Steril absorberende ikke-klæbende forbindelse, ca. 20 x 30 cm	5 stk.	0
B.3	Sterile absorbent non-adhesive surgical dressing bandage, approx. 20 x 30 cm	5 pcs.	0
B.4	Sterile kompresforbindinger, ca. 8 cm	10 ruller	0
B.4	Sterile compress bandages, approx. 8 cm	10 rolls	0
B.5	Imprægnerede net, ca. 10 x 10 cm, "Jelonet"	2 x 10 stk.	0
B.5	Impregnated nets, approx. 10 x 10 cm, "Jelonet"	2 x 10 pcs.	0
B.6	Steril absorberende forbindinger, ikke klæbende, ca. 10 x 10 cm	5 stk.	0
B.6	Sterile non-adhesive surgical dressing bandages, approx. 10 x 10 cm	5 pcs.	0
B.7	Gazeviskers, usterile	100 stk.	0
B.7	Gauze swaps, non-sterile	100 pcs.	0
B.8	Tubegaze til fingerforbindinger	10 stk.	0
B.8	Tube shaped gauze for finger bandaging	10 pcs.	0
B.9	Elastisk gazebind, ca. 4 cmx4m	10 ruller	0
B.9	Elastic gauze bandage, approx. 4 cmx4m	10 rolls	0
B.10	Hæfteplaster med sårpude, forskellige størrelser, lommepakning	1 pakke	0
B.10	Adhesive tape with compress, various sizes, pocket pack	1 pack	0
B.11	Hæfteplaster med sårpude, ca. 6 cm x 1m	5 pakker	0
B.11	Adhesive tape with compress, approx. 6 cm x 1m	5 packs	0
B.12	Fikseringsplaster Fixomull, ca. 10 cm x 10 m	1 rulle	0
B.12	Adhesive tape Fixomull, approx. 10 cm x 10 m	1 roll	0
B.13	Fikseringsplaster, "Englehud", ca. 2 cm	2 ruller	0
B.13	Adhesive tape, "peau d'ange", approx. 2 cm	2 rolls	0
B.18	Årepresse til massiv blødning	1 stk.	0
B.18	Tourniquet for massive bleeding	1 pc.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A2
C. Udstyr til hygiejne / Equipment for sanitation			
C.1	Engangshandsker af latex, usterile, X-large	100 stk.	0
C.1	Disposable gloves, latex, non-sterile, X-large	100 pcs.	0
C.2	Engangshandsker af latex, usterile, large	100 stk.	0
C.2	Disposable gloves, latex, non-sterile, large	100 pcs.	0
C.5	Vaskesvampe, éngangs	25 stk.	0
C.5	Bath sponge, disposable	25 pcs.	0
C.6	Beskyttelsesbriller	1 stk.	0
C.6	Protective goggles	1 pc.	0
C.7	Steril afdækning, ca. 50 x 50 cm	5 stk.	0
C.7	Sterile drape, approx. 50 x 50 cm	5 pcs.	0
C.8	Flydende medicinsk sæbe pH 4,5	250 ml	0
C.8	Liquid medical soap pH 4,5	250 ml	0
C.9	Mundbind 3 lags, bredde 18 cm	50 stk.	0
C.9	Mask, 3 layer, width 18 cm	50 pcs.	0
C.10	Ligpose voksen	1 stk.	0
C.10	Body bag adult	1 pc.	0
C.11	Engangshandsker af latex, usterile, medium	100 stk.	0
C.11	Disposable gloves, latex, non-sterile, medium	100 pcs.	0
C.12	Engangshandsker af latex, usterile, small	100 stk.	0
C.12	Disposable gloves, latex, non-sterile, medium	100 pcs	0
C.13	Engangsforklæde	10 stk.	0
C.13	Disposable apron	10 pcs.	0
C.14	Lagen, sterilt	1 stk	0
C.14	Bed sheet, sterile	1 pc.	0
C.15	Absorberende underlag, ca.6 lag, 60x40 cm, usteril, engangs	10 stk.	0
C.15	Absorbent underlay, approx.6 layers, 60x40 cm, non-sterile, disposable	10 pcs.	0
C.16	Opkastpose	10 stk.	
C.16	Vormit bag	10 pcs.	

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A2
D. Sårbehandling / Treatment of wounds			
D.1	Suturtape, "Steri-Strip™"	25 stk.	0
D.1	Suturetape, "Steri-Strip™"	25 pcs.	0
D.2	Klæbesutur, Dermabond	6 amuller	0
D.2	Adhesive suture, Dermabond	6 ampoules	0
D.4	Ikke-absorberbar sutur, 3.0; 45 cm. FS eller DS 25	12 stk.	0
D.4	Non-absorbable suture, 3.0; 45 cm. FS or DS 25	12 pcs.	0
D.5	Rustfri stålæske med låg, ca. 18x9x3 cm	1 stk.	0
D.5	Instrument box, stainless steel with lid, approx. 18x9x3 cm	1 pc.	0
D.6	Engangsskalpel, steril, nr. 23	5 stk.	0
D.6	Disposable scalpel, sterile, no. 23	5 pcs.	0
D.7	Kirurgisk saks med lige spids	1 stk.	0
D.7	Surgical scissors w/straight points	1 pc.	0
D.8	Kirurgisk saks med krum spids	1 stk.	0
D.8	Surgical scissors w/curved points	1 pc.	0
D.9	Kirurgisk pincet, ca. 14 - 16 cm	1 stk.	0
D.9	Surgical forceps (tweezers), approx. 14 -16 cm	1 pc.	0
D.10	Anatomisk pincet	1 stk.	0
D.10	Anatomical forceps (tweezers)	1 pc.	0
D.11	Splintpincet	1 stk.	0
D.11	Forceps, splinter (tweezers)	1 pc.	0
D.12	Pean, krum	2 stk.	0
D.12	Pean, curved	2 pcs.	0
D.13	Nåleholder	2 stk.	0
D.13	Suture - holder	2 pcs.	0
D.14	Barbermaskine (skraber), engangs	3 stk.	0
D.14	Safety razor, disposable single use	3 pcs.	0
D.15	Absorberbar hæmostatisk gelatinesvamp, ca. 1x1x1 cm	8 stk.	0
D.15	Haemostatic absorbable gelatine sponge, approx. 1x1x1 cm	8 pcs.	0
D.17	Ikke-absorberbar sutur, 5.0; 45 cm. FS eller DS25	12 stk.	0
D.17	Non-absorbable suture, 5.0; 45 cm. FS eller DS25	12 pcs.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A2
E. Udstyr til injektion og infusion / Equipment for injection and infusion			
E.1	Kanylebøtte	1 stk.	0
E.1	Box for contaminated needles	1 pc.	0
E.2	Afsprøjtningssvaps m. isopropylalkohol 70%	200 stk.	0
E.2	Alcohol swabs, isopropyl alcohol 70%	200 pcs.	0
E.3	Staserem	2 stk.	0
E.3	Tourniquet for i.v.	2 pcs.	0
E.4	Intravenøs kanyle, 1,1 x 32, 20 g (se note)	20 stk.	0
E.4	Intravenous cannula, 1.1 x 32, 20 g (see note)	20 pcs.	0
E.5	Fikseringsplaster til intravenøs kanyle	20 stk.	0
E.5	Fixing plaster for intravenous cannula	20 pcs.	0
E.6	Kanyle, steril til engangsbrug, 0,6 x 30 (~ 23G x 1 1/4) (se note)	50 stk.	0
E.6	Needle, sterile, disposablesingle use, 0.6 x 30 (~ 23G x 1 1/4) (see note)	50 pcs.	0
E.7	Sprøjter, sterile til engangsbrug, Luer, 2 ml	20 stk.	30 stk.
E.7	Syringes, sterile, single use, Luer, 2 ml	20 pcs.	30 pcs.
E.8	Sprøjter, sterile til engangsbrug, Luer, 5 ml	20 stk.	0
E.8	Syringes, sterile, single use, Luer, 5 ml	20 pcs.	0
E.9	Sprøjter, sterile til engangsbrug, Luer, 10 ml	20 stk.	0
E.9	Syringes, sterile, single use, Luer, 10 ml	20 pcs.	0
E.10	Infusionssæt med gevindtilslutning til E.4, sterile til engangsbrug	10 stk.	0
E.10	Infusion set with threaded connection for E.4, sterile, single use	10 pcs.	0
<p>E.4+E.6: Kanyle i kaliber 0,8 og intravenøs kanyle kaliber 1,2, som beskrevet i MFAG, er erstattet af kanyle i kaliber 0,6 og intravenøs kanyle kaliber 1,1.</p> <p>E.4+E.6: Needles size 0.8 and intravenous cannula size 1.2, as described in MFAG, are replaced with needles size 0.6 and intravenous cannula size 1.1</p>			
F. Udstyr til medicinsk undersøgelse / Equipment for medical examination			
F.1	Blodtryksmåler	1 stk.	0
F.1	Instrument for measuring blood pressure	1 pc.	0
F.2	Stetoskop	1 stk.	0
F.2	Stethoscope	1 pc.	0
F.3	Digitalt lægetermometer	2 stk.	0
F.3	Digital clinical thermometer	2 pcs.	0
F.4	Pencillygte m. batterier	2 stk.	0
F.4	Pencil lamp w/batteries	2 pcs.	0
F.5	Tungespatel, usterile, engangs	10 stk.	0
F.5	Tongue depressor, non-sterile, single use disposable	10 pcs.	0
F.7	Hypotermi Termometer	1 pc.	0
F.7	Hypotermi thermometer	1 stk.	0
<p>*)F.7 Termometret skal kunne måle minimum ned til 26°C. Bruges i endetarm.</p> <p>*)F.7 the thermomether have to be able to minimum measure down to 26°C. The thermomethre is used rectal.</p>			

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quentety A	MFAG A2
G. Udstyr til undersøgelse af blod og urin / Equipment for examination of blood and urine			
G.1	"Multistix® 5", urintest for blod/glukose/protein/nitrit/leukocytter	50 stk.	0
G.1	"Multistix® 5", Urinetest for blood/glucose/protein/nitrite/leukocytes	50 pcs.	0
G.2	Graviditetstest	2 stk.	0
G.2	Pregnancy test kit	2 pcs.	0
G.5	Malariatest, minimum for Plasmodium Falciparum, inkl. reagens og brugervejledning.	6 stk.	0
G.5	Malaria test, minimum for Plasmodium Falciparum, incl. reagens and userinstruction.	6 pcs.	0
G.6	Lancetter (fingerprikker)	6 stk.	0
G.6	Lancets (finger pricker)	6 pcs.	0
G.7	Udstyr til måling af blodsukker (se note)	0	0
G.7	Equipment for meusing bloodsugar (see note)	0	0
G.8	Hurtig test C-reaktiv protein (CRP)	5 stk.	0
G.8	Rapid test for C- reaktive proteine (CRP)	5 pcs.	0
G.2	Skal kun forefindes, HVIS der er kvinder om bord.		
G.2	Shall only be available IF women on board.		
G.5+G.6	Malariatest og Lancetter skal forefindes i skibe, der sejler til områder med malaria - uanset risiko.		
G.5+G.6	Malaria test and Lancets must be carried by all ships sailing to or in areas where malaria is recorded - irrespective of the risk.		
G.7	1 sæt skal medbringes af besætningsmedlemmer med insulinkriævende sukkersyge		
G.7	1 set to be carried by crew members with insulin dependent diabetes.		
H. Øjenundersøgelse og behandling / Eye examination and treatment			
H.1	Pencillygte med UV lys, inkl. batterier	1 stk.	0
H.1	Lamp with UV light w/batteries	1 pc.	0
H.2	Fluoretsstrips til farvning af øjet	10 stk.	0
H.2	Fluorescent colour for the eye	10 pcs.	0
H.3	Vatpinde, sterile	50 stk.	0
H.3	Cotton sticks, sterile	50 pcs.	0
H.4	Horners hulmejsel	1 stk.	0
H.4	Horners spud	1 pc.	0
H.5	Øjenklap	2 stk.	0
H.5	Eye patch	2 pcs.	0
H.6	Øjenskyllvæske, Natriumchlorid 0,9%, 250 ml	8 stk.	0
H.6	Eye irrigating liquid, Sodium chloride 0.9%, 250 ml	8 pcs.	0
I. Udstyr til tømning af urinblæren / Equipment for drainage of urinary bladder			
I.1	Engangskatetre m. krum spids, CH 14, til mænd	10 stk.	0
I.1	Catheters for single use w/ curved tip, CH 14 for men	10 pcs.	0
I.2	Urinkolbe, plastic	1 stk.	0
I.2	Bed bottle, plastic	1 pc.	0
I.3	Engangskatetre CH 14 til kvinder	5 stk.	0
I.3	Catheters for single use, CH 14 for women	5 pcs.	0

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quantity A	MFAG A2
J. Udstyr til forstuvninger og brud / Equipment for sprains and fractures			
J.1	ColdHot™ kulde/varme omslag, 11 x 27 cm	1 stk.	0
J.1	ColdHot™ pack 11x24 cm	1 pc.	0
J.2	Elastikbind, ca. 10 cmx5m	3 stk.	0
J.2	Elastic bandage, approx. 10 cmx5m	3 pcs.	0
J.3	Elastikbind, ca. 6 cmx5m	1 stk.	0
J.3	Elastic bandage, approx. 6 cmx5m	1 pc.	0
J.4	Trekantet tørklæde (Mitella)	2 stk.	0
J.4	Triangular scarf (Mitella)	2 pcs.	0
J.5	Sikkerhedsnåle	12 stk.	0
J.5	Safety pins	12 pcs.	0
J.6	Fingerskinne til immobilisering af ét led	5 stk.	0
J.6	Finger splint for immobilisation of one joint	5 pcs.	0
J.7	Fingerskinne til immobilisering af to led	5 stk.	0
J.7	Finger splint for immobilisation of two joints	5 pcs.	0
J.8	Polsterbind, ca. 10 cm x 2,7 m	10 stk.	0
J.8	Padded bandages, approx. 10 cm x 2.7 m	10 pcs.	0
J.9	Gipsbind, ca. 12 cmx2m	10 stk.	0
J.9	Plaster bandage, approx. 12 cmx2m	10 pcs.	0
J.10	Fikseringsplaster, Tensoplast, ca. 2 cmx2m	2 ruller	0
J.10	Adhesive tape, Tensoplast, approx. 2 cmx2m	2 rolls	0
K. Udstyr til immobilisering og transport / Equipment for immobilisation and transportation			
K.1	Båre, egnet til K.5 (se note)	1 stk.	0
K.1	Stretcher, suitable for K.5 (see note)	1 pc.	0
K.2	Oppustelig frakturskinne, senior, hel arm	1 stk.	0
K.2	Inflatable fracture splint, senior, full arm	1 pc.	0
K.3	Oppustelig frakturskinne, senior, hel ben	1 stk.	0
K.3	Inflatable fracture splint, senior, full leg	1 pc.	0
K.4	Stiv halskrave, indstillelig til forskellige positioner	2 stk.	0
K.4	Stiff neck collar, adjustable	2 pcs.	0
K.5	Langt rygbræt, inkl. seler (se note)	1 stk.	0
K.5	Spineboard, incl straps see note)	1 pc.	0
K.2 - K.4 pakkes og opbevares sammen. / K.2-K.4 packed and stored together			
K.1 + K.5 Meddelelser A kapitel IX B, bilag 3 / Notice A Chapter IX B, annex 3			

Grp. No.	Medicinsk udstyr / Medical equipment	Mængde/ Quentety A	MFAG A2
L. Tandlægeudstyr / Equipment for dental examination and treatment (Se note / see note)			
L.1	Coltoso®, færdigblandet tandcement i tuber à 8g (se note)	5 stk.	0
L.1	Coltoso®, dental filling material ready for use, tubes of 8 g (see note)	5 pcs.	0
L.2	Skyllekanyle	1 stk.	0
L.2	Dental rinse needle	1 pc.	0
L.3	Specialinstrument til tandcement	1 stk.	0
L.3	Special instrument for dental filling material	1 pc.	0
L.4	Mundspejl	1 stk.	0
L.4	Mouth mirror	1 pc.	0
L.5	Ekskavator	1 stk.	0
L.5	Excavator	1 pc.	0
L.6	Fyldningsfjerner	1 stk.	0
L.6	Filling remover	1 pc.	0
L.7	Rustfri stålæske med låg til tandudlægeudstyr	1 stk.	0
L.7	Instrument box, stainless steel with lid for dental equipment	1 pc.	0
<p>L: Tandlægeudstyr skal ikke forefindes på skibe med fart indenfor 175 sømil fra EU/EØS landes kyster. L.: Dental equipment shall not be on ships in voyage within 175 nautical miles from EU/EEA countries' cost L.1: Tandcement skal være inklusive instrument til uddrivning af cement fra tuben. L.1: Filling material shall include instrument for expulsion of the material from the tube.</p>			
M. Udstyr til sygepleje / Requisites for nursing			
M.1	Medicinmålebægre, ca. 25 ml, éngangs	20 stk.	0
M.1	Graduated measuring cup, approx. 25 ml, disposable	20 pcs.	0
M.2	Varmedunk	1 stk.	0
M.2	Hot-water bottle	1 pc.	0
M.3	Bækken	1 stk.	0
M.3	Bedpan	1 pc.	0
N. Insektbekæmpelsesmidler og kloreringsmidler til rensning af drikkevand/ Insecticides and chlorinating agents for purifying drinking water (se note / see note)			
N.1	Insektbekæmpelsesmiddel, pulver form	1 dåse	0
N.1	Insecticide, powder form	1 tin	0
N.2	Kloreringsmiddel til drikkevand, 100 g	2 stk.	0
N.2	Chlorinating agent for drinking water, 100 g	2 pcs.	0
N.3	Insektbekæmpelsesmiddel, væskeform	1 dåse	0
N.3	Insecticide, liquid form	1 tin	0
<p>N.1 + N.2 + N.3 Klor og insektbekæmpelsesmidler må ikke opbevares sammen med lægemidler og medicinsk udstyr. N.1 + N.2 + N.3 Chlor and insecticides must not be stored within the ship medicine chest</p>			
O. Fortegnelser, lægebog m.v. / Inventories, Medical Guide, etc.			
O.2	Lægebog for Søfarende, udgivet af Seahealth(se note)	1 stk.	0
O.2	Medical guide for seafarers issued by Seahealth (see Note)	1 pc.	0
O.3	Fortegnelse, kontroldokument og brugervejledning (se note)	1 stk kat. MFAG A2	0
O.3	Inventory, Control Document and User Instruction (see note)	1 pc. Cat.MFAG A2	0
O.5	Radio Medical Optegnelser	1 stk.	0
O.5	Radio Medical Records	1 pc.	0
O.6	MFAG (Medical First Aid Guide)	0	1 stk.
O.6	MFAG (Medical First Aid Guide)	0	1 pc.
<p>O.2 Skal også forefindes på engelsk, såfremt der om bord er påmønstret ikke dansktalende besætningsmedlemmer. / O.2 Must also be available in English, if crewmembers who don't speak Danish have signed on. O.3 skal være tilgængelig i digital eller printet udgave. O.3 must be available in digital or printed version.</p>			

Labels

<p>1.1 Gas / gas Medicinsk ilt / Medical Oxygen Indikation : Bevidstløse uden vejrtrækning, Bevidstløse med vejrtrækning, Medtaget almen tilstand. Slag mod kraniet, Større kvæstelser, Vejrtrækningsbesvær Indication : Unconscious, Weak general condition, Blows to the skull. Severe injuries, Difficulty breathing. Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring: Opbevares adskilt fra brandbare gasser og stoffer. Hold flasketemperaturen under 50°C / Storage: Keep away from flammable gases and substances. Keep bottle under 50°C</p>
<p>2.1 Inj. væske / inj. liquid Lidocain 10 mg/ml Indikation / Indication: Lokalbedøvelse / Local anaesthetic Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>2.2 Gel / gel Lidocain 2% Indikation/Indication: Lokalbedøvelse / Local anaesthetic Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>3.1 Tablet / tablet Paracetamol 500 mg Indikation / Indication: Svage smerter / Weak pain; MFAG tables 7, 8,13 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>3.2 Tablet / tablet Ibuprofen 400 mg Indikation / Indication: Moderate smerter / Moderate pain Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>3.4 Inj.væske / inj.liquid Δ Morfin 10 mg/ml Indikation / Indication: Stærke smerter / Severe pain; MFAG tables 7,8, 10,13 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>3.7 Gel / ointment Diclofenac 11.6 mg/g Indikation / Indication: Smerte / pain , inflammation I huden / inflammation of the skin. Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: < 30 C</p>
<p>3.8 Supp. / supp. Diclofenac 50 mg Indikation / Indication: Galdesten, nyresten, ledsmerter / Gall stone, kidney stone, pain in joint Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>3.9 tablet/tablets Morphine 10 mg Indikation / Indication: Stærke smerter / Severe pain; Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>4.1 Inj.væske / inj.liquid Adrenalin 1 mg/ml Indikation / Indication: Allergisk Shock / Allergic shock Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring: Læs indlgssedel/ Storage: Read packet insert</p>

<p>4.2 Kapsel / capsule Acrivastin 8 mg Indikation / Indication: Allergi / Allergy Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>4.3 Inj.væske / inj.liquid Hydrocortisonsuccinat 100 mg Indikation / Indication: Allergi / Allergy Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>4.5 Tablet / tablet Δ Cinnarizin 25 mg Indikation / Indication: Søsyge / Seasickness Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>4.6 Tablet / Tablets Prednisolon 25 mg Indikation / Indication Allergi / Allergy Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>5.1 Inj.væske / inj.liquid Naloxone 0,4 mg/ml Indikation/Indication: Morfin antidot / Morphine antidote MFAG tables 4, 13 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>5.3 Inj.væske / inj.liquid Atropin 1 mg/ml Indikation / Indication: Antidot / Antidote / MFAG table 17 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>5.4 Gel / gel Calcium gluconate 2 % Indikation / Indication: Antidot / Antidote / MFAG table 8, 16 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>5.5 Tablet / tablet Calciumlactogluconate 500 mg Indikation / Indication: Antidot / Antidote / MFAG tables 16 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>5.6 Afdelt pulver / powder Aktivt kul / Activated charcoal Indikation / Indication: Antidot / Antidote/ MFAG table 10 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>5.7 Væske / liquid Ethyl alcohol 99,5% Indikation/Indication: Antidot / Antidote/ MFAG table 19 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>6.2 Tablet / tablet Δ Diazepam 5 mg Indikation / Indication: Angst / Anxiety Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>

<p>6.3 Klyksma / enema ▲ Diazepam 4 mg/ml Indikation / Indication: Kramper / Convulsions; MFAG tables 4, 5, 6 Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>6.4 Inj.væske / inj liquid ▲ Haloperidol 5 mg/ml Indikation / Indication: Sindssyge / Mental disease Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>6.5 Tablet / tablet ▲ Zolpidem 10 mg Indikation / Indication: Søvnløshed / insomnia Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.1 Tablet / tablet Phenoxymethylpenicillin 660 - 800 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.2 Tablet / tablet Amoxicillin 500 mg Indikation / Indication: Infektioner / Infections / MFAG table 9 Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.6 Tablet / tablet Metronidazol 500 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.7 Suppositorie / suppository Metronidazol 500 mg Indikation / Indication: Infektioner / Infections / MFAG table 10 Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.8 Tablet / tablet Ciprofloxacin 250 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.9 Inj.substans. / inj.substance Ceftriaxon 1 g Indikation / Indication: Infektioner / Infections / MFAG table 10 Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.10 Vaccine / vaccine indeholdende / contain Tetanus Indikation / Indication: Forebyggelse mod stivkrampe / Tetanus prevention Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: 2- 8°C</p>
<p>7.11 Immunglobulin / immunglobuline Tetanus 250 I.E./ml / 250 I.U/ml Indikation / Indication: Forebyggelse mod stivkrampe / Tetanus prevention Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: 2- 8°C</p>

<p>7.12 Tablet / tablet Flucloxacillin 500 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.15 Tablet / Tablets Azithromycin 500 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.16 Tablet / Tablets Pivmecillinam 400 mg Indikation / Indication: Infektioner / Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>7.17 Tablet / Tablets Aciclovir 400 mg Indikation / Indication: virus-Infektioner / virus-Infections Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>8.1 Tablet / tablet Atovaquon/Proguanil 250/100 mg Indikation / Indication: Malaria forebyggelse / Malaria Prevention. Malaria behandling / Malaria treatment Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>8.2 Tablet / tablet Doxycyklin 100 mg Indikation / Indication: Malaria forebyggelse / Malaria prevention Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>8.4 tablet / tablet Artemether and lumefantrine 20 mg / 120 mg Indikation / Indication: Malaria behandling / Malaria treatment Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: ikke over 30° C / Not above 30° C</p>
<p>9.1 Spray / aerosol Glycerylnitrat / Glyceril trinitrate 0,4 mg/doses Indikation / Indication: Hjertekrampe / Angina pectoris Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: 2 - 25°C</p>
<p>9.3 Tablet / tablet Acetylsalicylsyre / acetylsalicylic acid 150 mg Indikation / Indication: Blodfortyndende / Blood thinning Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>9.4 Tablet / tablet Furosemid 40 mg Indikation / Indication: Vanddrivende / Diuretic Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>9.5 Inj.væske / inj.liquid Furosemid 10 mg/ ml Indikation / Indication: Vanddrivende / Diuretic MFAG tables 2, 9 Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory. Opbevaring / storage: normal temperatur / normal temperature</p>

<p>9.6 Tablet / tablet Atenolol 50 mg Indikation / Indication: Blodtryksforhøjelse / High blood pressure Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>9.8 Tablet / Tablets Apixaban 5 mg Indikation / Indication: Blodfortyndende / Blood diluting Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>9.9 inj. / inj. Tranexamsyre / Tranexamacid 100 mg/ml Indikation / Indication: Stopper blødninger / Stops Bleedings Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>10.2 Creme / lotion Miconazol 2% Indikation / Indication: Hudsvamp / Skinfungus Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: < 25°C</p>
<p>10.5 Creme / lotion Permethrin 5% Indikation / Indication: Fladlus og fnat / Crab lice and scabies Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>10.6 Shampoo / shampoo Permethrin 1% Indikation / Indication: Lus/ Lice Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>10.7 Creme / lotion Glukokortikoid gruppe II / glucocorticoid group II Indikation / Indication: Eksem / Eczema Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>10.8 Salve / ointment Fusidin 2 % Indikation / Indication: Hud infektion / Skin infection Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>11.1 Væske / liquid Chlorhexidin / Ethanol 0,5 - 5%/70 - 90% Indikation / Indication: Desinfektion af instrumenter / Disinfection of instruments Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>11.2 Væske / liquid Chlorhexidin 0,2% Indikation / Indication: Behandling af urene sår, mundskyl / Treatment of dirty wounds, mouth rinse. Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>12.6 Tablet / tablet Loperamid 2 mg Indikation / Indication: Diarré / Diarrhea Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>

<p>12.8 Kapsler / capsule Omeprazol 20 mg Indikation / Indication: Mavesår / Stomach ulcer Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>12.7 Pulver / powder Glucose / kalium / magnesium / natrium citrat/natrium/ Glucose/potassium/magnesium/sodium citrate/sodium 30/1,1/0,11/2,1 g Indikation / Indication: Væske- og salttab / Rehydration MFAG table 8, Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Opbevaring / storage: normal temperatur / normal temperature</p>
<p>12.9 Tablet eller oral opløsning / tablet or oral solution Antacida Indikation / Indication: Sure opstød og halsbrand / acid indigestion and heartburn Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>12.10 Rektalsalve / rectal ointment Antihæmorrhoid creme / Anti haemorrhoid ointment Indikation / Indication: Analkløe / Anal pruritus Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: < 25 C</p>
<p>12.13 Tablet / Tablets Ondasetron 4 mg Indikation / Indication: Kvalme og opkastning / Nausea and vomiting MFAG table 7, 8, 10, 13, 15, 20 Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / Storage normal temperatur.</p>
<p>12.14 Pulver / Power Movicol® Indikation / Indication: Forstoppelse / Constipation Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.2 Øjensalve / eyeointment Cinchocain 5 mg/g Indikation / Indication: Lokalbedøvelse af øje / Local anaesthetic of eye; MFAG table Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.3 Øjendråber / eyedrops Dexamethason 1 mg/ml Indikation / Indication: Allergi / Regnbuehindebetændelse / Allergic / Inflammatory eye diseases Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.4 Øjendråber / eyedrops Pilocarpin 20 mg/ml Indikation / Indication: Grøn stær / Glaucoma Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.6 Øjensalve / eyeointment Acyclovir 30 mg/g Indikation / Indication: Virusinfektion / Viral infection Ref: Se Fortegnelse brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>

<p>13.9 Væske / liquid Medium til at blødgøre ørevoks/ Medium to soften earwax Indikation / Indication: Blødgøring og løsning af ørevoks / Softening and dissolving of earwax Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.11 Øjendråber / eyedrops Tropicamid 1% Indikation / Indication: Øjeninflammation / Inflammation of eye Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.12 Øredråber / Eardrops Ciprofloxacin / Hydrocortison 10+2 mg Indikation: Øregangs eksem / Ear canal eczema Ref: Se Fortegnensens brugervejledning / Read user instruction in the inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>13.13 Øjensalve / eye ointment Chloramphenicol 1% Indikation: øjen infektion /Eye infection Ref: Se Fortegnensens brugervejledning / Read user instruction in the inventory. Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur</p>
<p>14.1 Kapsel / capsule Fluconazol 150 mg Indikation / Indication: Skedesvamp / Vaginal fungus Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>14.2 Tablet / tablet Methylergometrin 0,125 mg Indikation / Indication: Blødning efter fødsel / Bleeding after giving birth Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>15.1 Tablet / tablet Dextromethorphan 30 mg Indikation / Indication: Hoste / Cough Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>

<p>15.5 Næsespray / dråber / nose spray / drops Xylometazolin 1 mg/ml Indikation / Indication: Forkølelse / Common cold Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>15.7 Inhalations Spray / Inhalation Spray Salbutamol 0,1 mg Indikation / Indication: Astma / Astma Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>15.8 Inhalations spray / Inhalation spray Glucocorticoid 100-200 µg Indikation / Indication: Astma / Astma Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>16.1 Inf.væske / inf.liquid Natriumklorid / sodium chloride 9 mg/ml Indikation / Indication: Blodtab / Loss of blood; MFAg table 7 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>16.2 Inf.væske / inf.liquid Glucose og natrium / glucose and sodium 25/4,5 mg/ml Indikation / Indication: Væsketab / Rehydration Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>16.5 Inj. væske / inj.liquid Natriumklorid / sodium chloride 9 mg/ml Indikation / Indication: Væske til gennemskylning af intravenøs kanyle, sårbehandling, øjenbehandling / Fluid for flushing intravenous needle, treatment of wound, treatment of eyes Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>
<p>16.7 Væske / Liquid Sterilt vand / Sterile water Indikation / Indication: Væske til blanding af 7.9 / Fluid to mixing 7.9 Ref: Se Fortegnensens brugervejledning / Read the user instruction in the Inventory Søg samråd med Radio Medical / Consult Radio Medical Opbevaring / storage: normal temperatur / normal temperature</p>

INDEX

Abdominal infections	32; 33	Blodprop i hjertet, forebyggelse efter	43
Abstinenskrampe	28	Blodtab	62
Acetylsalicylic acid.	41	Blood diluting	44
Acetylsalicylsyre	41	Blood poisoning	34
Aciclovir	37; 55; 68; 71; 85; 86; 95; 97; 105	Blood-thinning	41
Acid indigestion	51	Blurred vision	54
Acrivastin	19	Blødning, abort	58
Acrivastine	19	Blødning, fødsel	58
Active carbon	26	Boil in the throat	32; 34
Acute diarrhoea	34	Bone infection	32
Acute mental disease	28	Bone-ache	14
Acute mental illness	29	Bones, infections	36
Acyclovir	55	Breathing difficulties	11
Adrenalin	18	Brud	28
Aktivt kul	26	Brugervejledning Lægemiddelstoffer	10
Akut diarré	33	Burning sensation, skin	46
Akut sindssyge	28; 29	Burns	15
Alcoholic withdrawal symptoms	28	Calciumgluconat	24
Alkoholabstinenser	28	Calciumgluconate	25
Allantoin	55	Calciumlactogluconat, calciumcarbonat.	25
Allergi	19	Calciumlactogluconate, calciumcarbonate.	25
Allergic asthma	19; 20	Cardiospasm	41
Allergic face oedema	19; 20	Caustic burns, fluorine	25
Allergic shock	18	Caustic burns, stomach, intestines	26
Allergisk ansigtsødem	19	Ceftriaxon	34
Allergisk astma	19	Ceftriaxone	34
Allergisk shock	18	Cerumen	55
Allergy	19; 20	Chlamydia infections	36
Amoxicillin.	31	Chlamydia infektioner	36
Amoxicilline	31	Chloramphenicol	57
Angina pectoris	40; 43	Chlorhexidin.	48
Angst	28; 29	Chlorhexidin/ethanol.	48
Antacida	51	Cholera	34
Antidote, morphine	22	Cinchocain.	53
Antihæmoride creme	51	Cinchocaine.	53
Anxiety	28; 29	Cinnarizin	20
Apixaban	44; 69; 86; 96; 105	Ciprofloxacin	33
Appendicitis	32; 33; 34	Ciprofloxacin	34
Arc eye	53	Cold	60
Artemether og lumefantrine	39	Constipation	52
Arthritis	15	Convulsions	28
asthma	60	Convulsions , universal	29
Asthma	61	Convulsions, poisoning	29
asthma-like breathing	61	Convulsions, Withdrawal	29
astma	60	Coronary thrombosis	15; 40; 41
Astma	43; 61	Coronary thrombosis, Preventiv treatment after	43
astmalignende vejrtrækning	61	Cough	59
Atenolol	43	Crab lice	45; 46
Atovaquon/Proguanil	38	Delusions	29
Atovaquone/Proguanil	38	Dexamethason.	53
Atropin	24	Dexamethasone.	54
Azithromycin	36; 68; 85; 95; 105	Dextromethorphan.	59
Back pains	13	Diarré	49
Bevidstløs med vejrtrækning	11	Diarrhoea	49; 50
Bevidstløs uden vejrtrækning	11	Diazepam	28; 29
Bihulebetændelse	31; 60	Diclofenac	15; 16
Bihulebetændelse	36	Difficulty falling asleep	30
Bleeding, after birth	59	Dirty wounds, Disinfection	48
Bleeding, miscarriage	59	Dislocations	28
Blindarmsbetændelse	32; 33; 34	Doxycycline.	39
Blodforgiftning	34	Doxycylin.	38
Blodfortyndende	44	Dysenteri	33
Blodfortynding	41	Dysentery	34
Blodprop i hjertet	14; 40; 41	Ear eczema	54

Eksem	46	Hands, Disinfection	48
Epilepsi	28	Hay fever	19
Epilepsy	29	Headache	13
Epinephrine	18	Heart pain	40; 41; 43
Ernæringserstatning	62	Heartburn	51
Ethyl alcohol	27	helvedesild	37
Ethyl alkohol	27	Helvedesild	55
Eye diseases, Allergic	54	Herpes	55
Eye diseases, Inflammatory	54	Herpes simplex	37
Eye infection	57	Herpes zoster	37
Eyes, Treatment	63	High blood pressure	42; 43
Feber	13	Hjertekrampe	40; 41
Febertilstande	49	Hostestillende	59
Female reproductions organs, infection	33	Hovedpine	13
Fever	13	Hudinfektioner	47
Feverish conditions	50	Hudkløe	46
Fladlus	45; 46	Hudsvamp	45; 58
Flucloxacillin	36	Hudsvie	46
Fluconazol	58	Hydrocortisonsuccinat.	19
Fluconazole.	58	Hydrocortisonsuccinate	20
Flushing, intravenous needle	63	Hypoglycaemic coma	23
Fnat	45; 46	Hæmorider	51
Food substitute	63	Hænder, desinficering	48
Forbrændinger	14	Høfeber	19
Forgiftning, insektgift	24	Ibuprofen.	13; 14
Forgiftninger med methanol, ethylene glycol	27	Indsovningsbesvær	30
Forgiftninger, mave, tarm	26	Infection of female reproductions organs	32
Forhøjet blodtryk	41; 42; 43	Infections in connection with ulcer	36
Forkølelse	60	Infections of the oral cavity	36
Forstoppelse	52	Infections, oral cavity	31
Fortegnelse og kontrol dokument over lægemiddelstoffer	65	Infektioner, stafylokokker	36
		Inflammation i huden	15
Fortegnelse og Kontrol dokument over medicinsk udstyr	74	Inflammation of the middle ear	36
		Inflammation of the skin	15
Fractures	14; 15; 28	Inflammation of the throat	36
Frontal sinusitis	60	Inflammation, middle ear	31
Fungus	45	Inflammation, throat	31
Furosemid	41; 42	Inflammatoriske øjensygdomme	56
Furosemide	42	Inflammatory eye diseases	56
Fusidin	47; 69; 86; 96; 97; 105	Injuries	15
Galdesten	14	Insect bites	46
Gall stone	15	Insecticides	24
Gennemskylning, intravenøskanyle	63	Insektbid	46
Gigt	15	Insektgifte	24
Glaucoma	54	Insomnia	30
Glucagon	23	Instrumenter, desinficering	48
Glucocorticoid til inhalation	61	Instruments, Disinfection	48
Glucocorticoid group II	46	Insulin shock	23
Glucose and sodium chloride.	63	Insulinshock	23
Glucose og natriumklorid.	62	Insulintilfælde	23
Glukokortikoid gruppe II	46	Intact skin, Disinfection	48
Glyceryl trinitrate	40	Intakt hud, desinficering	48
Glycerylnitrat	40	Intestinal infections	32; 33
Glycocorticoid for inhalation	61	Inventory and Control Document	65
Gonoré	33	Inventory and Control document of medical equipment	74
Gonorrhoea	34	Itching, rectum	51
Grøn stær:	54	Kidney stone	15
Haemorrhoids	51	Kloramfenikol	57
Hallucinationer.	29	Kløe, endetarmen	51
Hallucinations.	29	Kløestillende	12
Haloperidol	29	Knoglebrud	13
Haloperidole.	29	Knogleinfektion	32; 36
Halsbetændelse	31; 36	Knoglesmerter	13
Halsbrand	51	Kolera	33
Halsbyld	32; 33	Kramper	28

Kramper, forgiftninger	28	MFAG table 16	25
Kramper, universelle	28	MFAG table 17	24
Kvalme	52	MFAG table 19	27
Kvalme og opkastninger	29	MFAG table 2, 3, 5, 9, 11	11
Kvæstelser	14	MFAG table 2, 9	42
Kæbehulebetændelse	60	MFAG table 4, 13	22
Labels	83	MFAG table 4, 5, 6	28
Lavt blodsukker	23	MFAG table 7	53
Ledskred	28	MFAG table 8, 10, 11:	50
Ledsmerter	13	MFAG table 8, 16	25
Legionella	36	MFAG table 9	36
Lice	45; 46	Miconazol	45
Lidocain	11; 12	Miconazole	45
Lidocaine	12	Moderate pains:	14
	104	Moderate smerter:	13
	104	Modgift, morfin	22
Local anaesthetic, eye	53	Morfin	14
Local anaesthetics	12	Morfin Δ	16
Lokalbedøvelse	11; 12	Morphine	15
Lokalbedøvelse, øjet	53	Morphine Δ	17
Loperamid	49	Motoric restlessness,	29
Loperamide.	49	Motorisk uro	29
Loss of blood	62	Mouth rinse	48
Loss of fluid	50	Movicol®	52
Loss of liquid	62	Mundhuleinfektioner	31; 36
Loss of liquids	63	Mundskylning	48
Low blood sugar	23	Muscle pains	14
Lung edema	42	Muskelsmerter	13
Lungebetændelse	31; 34; 36	Nail fungus	58
Lungebetændelse, giftige kemiske stoffer	31	Naloxon	22
Lungepest	33	Naloxone	22
Lus	45; 46	Natriumklorid	63
Major wounds	15; 34	Natriumklorid.	62
Malaria, forebyggelse	38	Nausea	52
Malaria, prevention	38; 39	Nausea and vomiting	29
Malarone®	38	Neglesvamp	58
Mani, vildskab	29	Nyrebækkenbetændelse	33; 34
Mania, wildness	29	Nyresten	14
Mavesår	50	Nældefeber	19
Mavesårsinfektioner	36	Omeprazol	50
Maxillary sinusitis	60	Ondansetron	52
Medical oxygen	11	Open fractures	34
Medicinsk ilt.	11	Opkastninge	52
Medtaget almen tilstand	11	Opkastninger	49
Mellemørebetændelse	31; 36; 60	Otitis media	60
Meningitis	34	Oxygen	11
Menstruationssmerter	13	P. Falciparum malaria	39
Mental restlessness	29	P. Falciperum malaria	39
Methylergometrin.	58	Pain	15
Methylergometrine.	59	Pain during defecation	51
Metronidazol.	32; 33	Pain in joints	14
Metronidazole	32	Pains, eye	54
MFAG tabel 4, 5, 6	29	Pandehulebetændelse	60
MFAG tabel 8, 10, 11	49	Paracetamol	13
MFAG tabel 10	26; 32; 34	Paracetamol.	13
MFAG tabel 16	25	Period pains	14
MFAG tabel 17	24	Permethrin.	45; 46
MFAG tabel 19	27	Phenoxymethylpenicillin	31
MFAG tabel 4, 13	22	Phenoxymethylpenicillin	31
MFAG tabel 7	53; 62	Pilocarpin	54
MFAG tabel 7, 8,10,13	14	Pilocarpine	54
MFAG tabel 7,8,10,13	15	Pivmecillinam	37; 68; 85; 95; 105
MFAG tabel 8, 16	24	Pneumonia	31; 34; 36
MFAG tabel 9	31; 36	Pneumonia, poisonous chemical substances	31
MFAG table 10	26; 34	Pneumonic plague	34

Poisoning, insecticides	24	Søvnløshed	30
Poisoning, methanole, ethylene glycole	27	Sårbehandling.	63
Poisoning, stomach, intestines	26	Tandbyld	32; 33
Prednisolon	21	Tarminfektioner	32
Prenisolon	21	Tetanus holdig vaccine	35
Psoriasis	46	Tetanus vaccine	35
Psykisk uro	29	tetanustoxid	35
Pyelitis	34	Tooth abscess	32; 34
Rainbow vision	54	Tranexamacid	44
Regnbuesyn	54	Tranexamsyre	44
Relieves itching	12	Tropicamid	56
restlessness	28	Tågesyn	54
Rygsmarter	13	Udsættende vejtrækning	22
Salbutamol	60	Unconscious with breathing	11
Salbutamol.	60	Unconscious without breathing	11
Salttab	49	Underlivsinfektioner	32
Scabies	45; 46	Urene sår, desinficering	48
Seasickness	20	Urethritis	34
Sepsis	34	Urinary tract infection	37
Severe bronchitis	34	Urinrørsbetændelse	33
Severe injuries	11	Urinvejsinfektioner	37
Severe pain	15; 17	uroilstande	28
Shingles	55	Urticaria	19
Shock, forebyggelse	62	User instruction Activ Phamaceutical Ingredients	10
Shock, prevention	62	Vaginal fungus	58
Sinusitis	31; 36; 60	Vand i lungerne	41; 42
Skin fungus	45; 58	Vand til opløsning af 7.9 Ceftriaxon pulver	64
Skininfections	47	Variceller	37
Skoldkopper	37	Vejtrækningsbesvær	11
Smerter	15	Vomiting	50; 52
Smerter i hjertet	40; 41	Vrangforestillinger	29
Smerter, afføring	51	Væsketab	49; 62
Smerter, øje	54	Water for dissolution of 7.9 Ceftriaxon powder	64
Sodium chloride	62; 63	Weak general condition	11
Solskoldning	46	Weak pains:	13
Staphylococci	36	Welders eye	53
Sterile water	64	Wounds, Treatment	63
Sterilt vand	64	Xylometazolin.	60
Stivkrampevaccine	35	Xylometazoline.	60
Stomach ulcer	50	Zolpidem	30
Stopper blødninger	44	Ændringer /Changes	107
Stops Bleedings	44	Ætsninger, fluorholdige	24; 25
Stops breathing	22	Ætsninger, mave, tarm	26
Store sårskader	14; 34	Øjenbehandling	63
Stærke smerter	16	Øjenbetændelse	57
Stærke smerter:	14	Øjensygdom, Allergiske	53
Større kvæstelser	11	Øjensygdom, Inflammatoriske	53
Sunburns	46	Øregangseksem	53
Sure opstød	51	Ørevoks.	55
Suture of wounds	12	Åbne knoglebrud	34
Svage smerter	13		
Svamp	45		
Svamp i skeden	58		
Svejseøjne	53		
Svær bronkitis	33		
Syning af sår	11		
Søsyge	20		

Rekvisition lægemiddelstoffer og medicinsk udstyr

**Requisition Active Pharmaceutical Ingredient and
medical equipment**

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
1.1	V03AN01	Medicinsk ilt	Gas	
1.1	V03AN01	Oxygen, Medical	Gas	
1.3	-	Ekstra iltflaske	Gas	
1.3	-	Spare oxygen cylinder	Gas	
2.1	N01BB02	Lidocain	Inj. Væske	
2.1	N01BB02	Lidocaine	Inj. Liquid	
2.2	D04AB01	Lidocain	Gel	
2.2	D04AB01	Lidocaine	Gel	
3.1	N02BE01	Paracetamol	Tablet	
3.1	N02BE01	Paracetamol	Tablets	
3.2	M01AE01	Ibuprofen	Tablet	
3.2	M01AE01	Ibuprofen	Tablets	
3.4	N02AA01	Morfin Δ	Inj. Væske	
3.4	N02AA01	Morphine Δ	Inj. Liquid	
3.7	M02AA15	Diclofenac	Gel	
3.7	M02AA15	Diclofenac	Gel	
3.8	M01AB05	Diclofenac	Supp.	
3.8	M01AB05	Diclofenac	Supp	
3.9	N02AA01	Morfin Δ	Tablet	
3.9	N02AA01	Morphine Δ	Tablets	
4.1	C01CA24	Adrenalin	inj. Væske	
4.1	C01CA24	Epinephrine	inj. Liquid	
4.2	R06AX18	Acrivastine	Kapsel	
4.2	R06AX18	Acrivastine	Capsula	
4.3	H02AB09	Hydrocortison	inj. tørstof + væske	
4.3	H02AB09	Hydrocortisonsuccinate	inj. Subst + solvens	
4.5	N07CA02	Cinnarizin Δ	Tablet	
4.5	N07CA02	Cinnarizine Δ	Tablets	
4.6	H02AB06	Prednisolon	Tablet	
4.6	H02AB06	Prednisolon	Tablets	
5.1	V03AB15	Naloxon	Inj. Væske	
5.1	V03AB15	Naloxon	Inj. Liquid	
5.2	H04AA01	Glucagon "K"	inj. tørstof + væske	
5.2	H04AA01	Glucagon "R"	inj. Subst + solvens	
5.3	A03BA01	Atropine	Inj. Væske	
5.3	A03BA01	Atropine	Inj. Liquid	
5.4	D11AX03	Calcium gluconate	gel	
5.4	D11AX03	Calcium gluconate	gel	
5.5	A12AA02 / A12AA06	Calcium lactogluconate	Brusetablet	
5.5	A12AA02 / A12AA06	Calcium lactogluconate	Effervescent tablet	
5.6	A07BA01	Aktivt kul	pulver	
5.6	A07BA01	Charcoal, activated	powder	
5.7	-	Ethyl alcohol	Væske	
5.7	-	Ethyl alcohol	Liquid	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
6.2	N05BA01	Diazepam Δ	Tablet	
6.2	N05BA01	Diazepam Δ	Tablets	
6.3	N05BA01	Diazepam Δ	Klysma	
6.3	N05BA01	Diazepam Δ	Enema	
6.4	N05AD01	Haloperidol Δ	Inj. Væske	
6.4	N05AD01	Haloperidol Δ	Inj. Liquid	
6.5	N05CF02	Zolpidem Δ	Tablet	
6.5	N05CF02	Zolpidem Δ	Tablets	
7.1	J01CE02	Phenoxymethylpenicillin	Tablet	
7.1	J01CE02	Phenoxymethylpenicillin	Tablets	
7.2	J01CA04	Amoxicillin	Tablet	
7.2	J01CA04	Amoxicillin	Tablets	
7.6	P01AB01	Metronidazol	Tablet	
7.6	P01AB01	Metronidazol	Tablets	
7.7	P01AB01	Metronidazol	Supp.	
7.7	P01AB01	Metronidazol	Supp.	
7.8	J01MA02	Ciprofloxacin	Tablet	
7.8	J01MA02	Ciprofloxacin	Tablets	
7.9	J01DD04	Ceftriaxon	inj. Tørstof	
7.9	J01DD04	Ceftriaxon	inj. Subst	
7.10	-	Tetanus holdig vaccine "K"	Inj. Væske	
7.10	-	Vaccine containing Tetanus "R"	Inj. Liquid	
7.11	J06BB02	Tetanus immunglobulin "K"	Inj. Væske	
7.11	J06BB02	Tetanus immunglobulin "R"	Inj. Liquid	
7.12	J01CF05	Flucloxacillin	Tablet	
7.12	J01CF05	Flucloxacillin	Tablets	
7.15	J01FA10	Azithromycin	Tablet	
7.15	J01FA10	Azithromycin	Tablets	
7.16	J01CA08	Pivmecillinam	Tablet	
7.16	J01CA08	Pivmecillinam	Tablets	
7.17	J05AB01	Aciclovir	Tablet	
7.17	J05AB01	Acyclovir	Tablets	
8.1	P01BB51	Atovaquon/Proguanil	Tablet	
8.1	P01BB51	Atovaquon/Proguanil	Tablets	
8.2	J01AA02	Doxycycline	Tablet	
8.2	J01AA02	Doxycycline	Tablets	
8.4	P01BF01	Artemether and lumefantrine	Tablets	
8.4	P01BF01	Artemether and lumefantrine	Tablets	
9.1	C01DA02	Glyceryl nitrat	Mund pray	
9.1	C01DA02	Glyceryl trinitrate	Oral spray	
9.3	B01AC06	Acetylsalisylsyre	Tablet	
9.3	B01AC06	Acetylsalicylic Acid	Tablets	
9.4	C03CA01	Furosemid	Tablet	
9.4	C03CA01	Furosemide	Tablets	
9.5	C03CA01	Furosemide	Inj. væske	
9.5	C03CA01	Furosemide	Inj. Liquid	
9.6	C07AB03	Atenolol	Tablet	
9.6	C07AB03	Atenolol	Tablets	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
9.8(π)	B01AF02	Apixaban	Tablet	
9.8(π)	B01AF02	Apixaban	Tablets	
9.9(π)	B02AA02	Tranexamsyre	Inj. væske	
9.9(π)	B02AA02	Tranexamsyre	Inj. liquid	
10.2	D01AC02	Miconazol	Creme	
10.2	D01AC02	Miconazol	Lotion / Cream	
10.5	P03AC04	Permethrin	Creme	
10.5	P03AC04	Permethrin	Lotion / Cream	
10.6	P03AC04	Permethrin	Shampoo	
10.6	P03AC04	Permethrin	Shampoo	
10.7	D07AB02	Glukocorticoid group II	Creme	
10.7	D07AB02	Glukocorticoid group II	Lotion / Cream	
10.8	D06AX01	Fusidin	Cream / Salve	
10.8	D06AX01	Fusidine	Cream / Lotion	
11.1	-	Chlorhexidine/ethanol	Væske	
11.1	-	Chlorhexidine/ethanol	Liquid	
11.2	-	Chlorhexidine	Væske	
11.2	-	Chlorhexidine	Liquid	
12.6	A07DA03	Loperamid	Tablet	
12.6	A07DA03	Loperamid	Tablet	
12.7	-	Glucose/potassium/magnesium /sodium	Powder	
12.7	-	Glucose/potassium/magnesium /sodium	Powder	
12.8	A02BC01	Omeprazol	Tablet / Kapsel	
12.8	A02BC01	Omeprazol	Tablet / Capsula	
12.9	A02A	Antacida	Tablet eller oral opløsning	
12.9	A02A	Antacid	Tablet or oral solution	
12.10	C05A	Hæmoride creme(skal indeholde glukokortikoid and smertestillende)	Creme/Salve	
12.10	C05A	Anti haemorrhoid agent (Must contain glucocorticoid and analgesi)	Lotion/ointment	
12.13	A04AA01	Ondansetron	Tablet	
12.13	A04AA01	Ondansetron	Tablets	
12.14	A06AD65	Movicol®	Pulver	
12.14	A06AD65	Movicol®	Powder	

Grp. No.	ATC-code	Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
13.2	S01HA06	Cinchocain	Salve	
13.2	S01HA06	Cinchocaine	Ointment	
13.3	S01BA01	Dexamethason	Dråber	
13.3	S01BA01	Dexamethasone	Drops	
13.4	S01EB51	Pilocarpin	Dråber	
13.4	S01EB51	Pilocarpine	Drops	
13.6	S01AD03	Aciclovir	Salve	
13.6	S01AD03	Acyclovir	Ointment	
13.8	S01AA13	Fusidin	Dråber / gel	
13.8	S01AA13	Fusidin	Drops / gel	
13.9	-	Medium til at blødgøre ørevoks, inkl. Øresprøjte	-	
13.9	-	Medium to soften earwax, incl. Earsyringe	-	
13.11	S01FA06	Tropicamid	Dråber	
13.11	S01FA06	Tropicamide	Drops	
13.12	S02CA03	Ciprofloxacin / Hydrocortison	Øredråber	
13.12	S02CA03	Ciprofloxacin / Hydrocortisone	Eardrops	
13.13(π)	S01AA01	Kloramfenikol	øjensalve	
13.13(π)	S01AA01	Chloramphenicol	eye ointment	
14.1	J02AC01	Fluconazol	Kapsel	
14.1	J02AC01	Fluconazole	Capsula	
14.2	G02AB01	Methylergometrin	Tablet	
14.2	G02AB01	Methylergometrine	Tablets	
15.1	R05DA09	Dextromethorphan	Tablet	
15.1	R05DA09	Dextromethorphan	Tablets	
15.4	R03BA02	Budesonid	Turbohaler	
15.4	R03BA02	Budesonide	Turbohaler	
15.5	R01AA07	Xylometazolin	Dråber / spray	
15.5	R01AA07	Xylometazoline	Drops / spray	
15.7	R03AC03	Salbutamol	Spray	
15.7	R03AC03	Salbutamol	Spray	
15.8	R03BA01	Glucocorticoid (inhalation)	Spray	
15.8	R03BA01	Glucocorticoid (inhalation)	Spray	
16.1	B05BB01	Natriumklorid	Inf. Væske	
16.1	B05BB01	Sodium chloride	Inf. Liquid	
16.2	B05BB02	Glucose + Natriumklorid	Inf. Væske	
16.2	B05BB02	Glucose + sodiumchlorid	Inf. Liquid	
16.5	V07AB	Natriumklorid 20 ml	Væske	
16.5	V07AB	Sodium chloride 20 ml	Liquid	
16.7	V07AB	Sterilt vand til blanding af 7.9	Væske	
16.7	V07AB	Sterile water to mix 7.9	Kiquid	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
A.1	Mobilt genoplivningsudstyr, klargjort	
A.1	Mobile resuscitation equipment, ready for use	
A.4	Tungeholdere, størrelse 2	
A.4	Guidel airway, size 2	
A.5	Tungeholdere, størrelse 3	
A.5	Guidel airway, size 3	
A.6	Tungeholdere, størrelse 4	
A.6	Guidel airway, size 4	
A.7	Ekstra ventilationspose med ventil og voksenmaske	
A.7	Extra ventilation bag with valve and mask	
A.8	Kraftig klædesaks	
A.8	Strong scissors for clothes	
A.9	Sugekatetre, lige, CH 12	
A.9	Suction catheters, straight, CH 12	
A.10	Tæppe til behandling af hypotermi	
A.10	Blanket for treatment of hypothermia	
A.11	Almindelige tæpper	
A.11	Ordinary blankets	
A.12	Saturationsmåler (fingermodel)	
A.12	Puls oximeter (finger)	
A.13	Hudson maske med reservoirpose med ca. 1,8 m slange	
A.13	Hudson mask with reservoir bag with approx. 1.8 m tube	
A.18	Hjertestarter (AED)	
A.18	Automatic extern difibrillator (AED)	
A.19	Iltbrille	
A.19	Nasal Cannula	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
B.1	Sterile gazeekompresser, ca.5x5 cm, ca.8 lag	
B.1	Sterile gauze compress, approx.5x5 cm, approx. 8 layers	
B.2	Sterile gazeekompresser, ca. 10 x 10 cm, ca. 8 lag	
B.2	Sterile gauze compress, approx.10x10 cm, approx. 8 layers	
B.3	Steril absorberende ikke-klæbende forbindelse, ca. 20 x 30 cm	
B.3	Sterile absorbent non-adhesive surgical dressingbandage, approx. 20 x 30 cm	
B.4	Sterile kompresforbindinger, ca. 8 cm	
B.4	Sterile compress bandages, approx. 8 cm	
B.5	Imprægnerede net, ca. 10 x 10 cm, Jelonet	
B.5	Impregnated nets, approx. 10 x 10 cm, Jelonet	
B.6	Sterile absorberende forbindinger, ikke klæbende, ca. 10 x 10 cm	
B.6	Sterile absorbent bandages, non-adhesive surgical dressingbandages, approx. 10 x 10 cm	
B.7	Gazeviskers, usterile	
B.7	Gauze swaps, non-sterile	
B.8	Tubegaze til fingerforbindinger	
B.8	Tube shaped gauze for finger bandaging	
B.9	Elastisk gazebind, ca. 4 cmx4m	
B.9	Elastic gauze bandage, approx. 4 cmx4m	
B.10	Hæfteplaster med sårpude, forskellige størrelser, lommepakning	
B.10	Adhesive tape with compress, various sizes, pocket pack	
B.11	Hæfteplaster med sårpude, ca. 6 cmx1m	
B.11	Adhesive tape with compress, approx. 6 cmx1m	
B.12	Fikseringsplaster Fixomull, ca. 10 cm x 10 m	
B.12	Adhesive tape Fixomull, approx. 10 cm x 10 m	
B.13	Fikseringsplaster, "Englehud", ca. 2 cm	
B.13	Adhesive tape, "peau d'ange", approx. 2 cm	
B.18	Årepresse til massigv blødning	
B.18	Tourniquet for massive bleeding	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
C.1	Engangshandsker af latex, usterile, X-large	
C.1	Disposable gloves, latex, non-sterile, X-large	
C.2	Engangshandsker af latex, usterile, large	
C.2	Disposable gloves, latex, non-sterile, large	
C.5	Vaskesvampe, éngangs	
C.5	Bath sponge, disposable	
C.6	Beskyttelsesbriller	
C.6	Protective goggles	
C.7	Steril afdækning, ca. 50 x 50 cm	
C.7	Sterile drape, approx. 50 x 50 cm	
C.8	Flydende medicinsk sæbe pH 4,5	
C.8	Liquid medical soap pH 4,5	
C.9	Mundbind 3 lags, bredde 18 cm	
C.9	Mask, 3 layer, width 18 cm	
C.10	Ligpose voksen	
C.10	Body bag adult	
C.11	Engangshandsker af latex, usterile, medium	
C.11	Disposable gloves, latex, non-sterile, medium	
C.12	Engangshandsker af latex, usterile, small	
C.12	Disposable gloves, latex, non-sterile, small	
C.13	Engangsforklæde	
C.13	Disposable apron	
C.14	Lagen, sterilt	
C.14	Bed sheet, sterile	
C.15	Absorberende underlag, ca.6 lag, 60x40 cm, usteril, engangs	
C.15	Absorbent underlay, approx.6 layers, 60x40 cm, non-sterile, disposable	
C.16	Opkastpose	
C.16	Vomit bag	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
D.1	Suturtape, Steri-Strip™	
D.1	Suturtape, Steri-Strip™	
D.2	Klæbesutur, Dermabond	
D.2	Adhesive suture, Dermabond	
D.4	Ikke-absorberbar sutur, 3.0; 45 cm. FS eller DS 25	
D.4	Non-absorbable suture, 3.0; 45 cm. FS or DS 25	
D.5	Rustfri stålæske med låg, ca. 18x9x3 cm	
D.5	Instrument box, stainless steel with lid, approx. 18x9x3 cm	
D.6	Engangsskalpel, steril, nr. 23	
D.6	Disposable scalpel, sterile, no. 23	
D.7	Kirurgisk saks med lige spids	
D.7	Surgical scissors w/straight points	
D.8	Kirurgisk saks med krum spids	
D.8	Surgical scissors w/curved points	
D.9	Kirurgisk pincet, ca. 14 - 16 cm	
D.9	Surgical forceps (tweezers), approx. 14 -16 cm	
D.10	Anatomisk pincet	
D.10	Anatomical forceps (tweezers)	
D.11	Splintpincet	
D.11	Forceps, splinter (tweezers)	
D.12	Pean, krum	
D.12	Pean, curved	
D.13	Nåleholder	
D.13	Suture - holder	
D.14	Barbermaskine (skraber), engangs	
D.14	Safety razor, disposable single use	
D.15	Absorberbar hæmostatisk gelatinesvamp, ca.1x1x1 cm	
D.15	Haemostatic absorbable gelatine sponge, approx.1x1x1 cm	
D.17	Ikke-absorberbar sutur, 5.0; 45 cm. FS eller DS25	
D.17	Non-absorbable suture, 5.0; 45 cm. FS eller DS25	
E.1	Kanylebøtte	
E.1	Box for contaminated needles	
E.2	Afsprøjtningssvaps m. isopropylalkohol 70 %	
E.2	Alcohol swabs, isopropyl alcohol 70 %	
E.3	Staserem	
E.3	Tourniquet for i.v.	
E.4	Intravenøs kanyle, 1,1 x 32, 20 g	
E.4	Intravenous needle, 1.1 x 32, 20 g	
E.5	Fikseringsplaster til intravenøs kanyle	
E.5	Fixing plaster for intravenous needle	
E.6	Kanyle, steril til engangsbrug, 0,6 x 30 (~ 23G x 1 1/4)	
E.6	Needle, sterile, disposable single use, 0.6 x 30 (~ 23G x 1 1/4)	
E.7	Sprøjter, sterile til engangsbrug, Luer, 2 ml	
E.7	Syringes, sterile, single use, Luer, 2 ml	
E.8	Sprøjter, sterile til engangsbrug, Luer, 5 ml	
E.8	Syringes, sterile, single use, Luer, 5 ml	
E.9	Sprøjter, sterile til engangsbrug, Luer, 10 ml	
E.9	Syringes, sterile, single use, Luer, 10 ml	
E.10	Infusionsæt med gevindtilslutning til E.4, sterile til engangsbrug	
E.10	Infusion set with threaded connection for E.4, sterile, single use	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
F.1	Blodtryksmåler	
F.1	Instrument for measuring blood pressure	
F.2	Stetoskop	
F.2	Stethoscope	
F.3	Digitalt lægetermometer	
F.3	Digital clinical thermometer	
F.4	Pencillygte m. batterier	
F.4	Pencil lamp w/batteries	
F.5	Tungespatel, usterile, engangs	
F.5	Tongue depressor, non-sterile, single use disposable	
F.7	Hypotermi Termometer	
F.7	Hypotermi thermometer	
G.1	Multistix® 5, urintest for blod/glukose/protein/nitrit/leukocytter	
G.1	Multistix® 5, Urinetest for blood/glucose/protein/nitrite/leukocytes	
G.2	Graviditetstest	
G.2	Pregnancy test kit	
G.5	Malariatest, minimum for Plasmodium Falciparum, inkl. reagens og brugervejledning.	
G.5	Malaria test, minimum for Plasmodium Falciparum, incl. reagens and userinstruction.	
G.6	Lancetter (fingerprikker)	
G.6	Lancets (finger pricker)	
G.7	Udstyr til måling af blodsukker	
G.7	Equipment for measuring bloodsugar	
G.8	Hurtig test C-reaktiv protein (CRP)	
G.8	Rapid test for C- reaktive proteine (CRP)	
H.1	Pencillygte med UV lys, inkl. batterier	
H.1	Lamp with UV light w/batteries	
H.2	Fluoretsstrips til farvning af øjet	
H.2	Fluorescent colour for the eye	
H.3	Vatpinde, sterile	
H.3	Cotton sticks, sterile	
H.4	Horners hulmejsel	
H.4	Horners spud	
H.5	Øjenklap	
H.5	Eye patch	
H.6	Øjenskylllevæske, Natriumchlorid 0,9%, 250 ml	
H.6	Eye irrigating liquid, Sodium chlorid e 0.9%, 250 ml	
I.1	Engangskatetre m. krum spids, CH 14, til mænd	
I.1	Catheters for single use w/ curved tip, CH 14 for men	
I.2	Urinkolbe, plastic	
I.2	Bed bottle, plastic	
I.3	Engangskatetre CH 14 til kvinder	
I.3	Catheters for single use, CH 14 for women	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
J.1	ColdHot™ kulde/varme omslag, ca.11 x 27 cm	
J.1	ColdHot™ pack approx. 11 x 27 cm	
J.2	Elastikbind, ca. 10 cm x 5m	
J.2	Elastic bandage, approx. 10 cm x 5m	
J.3	Elastikbind, ca. 6 cm x 5m	
J.3	Elastic bandage, approx. 6 cm x 5m	
J.4	Trekantet tørklæde (Mitella)	
J.4	Triangular scarf (Mitella)	
J.5	Sikkerhedsnåle	
J.5	Safety pins	
J.6	Fingerskinne til immobilisering af ét led	
J.6	Finger splint for immobilisation of one joint	
J.7	Fingerskinne til immobilisering af to led	
J.7	Finger splint for immobilisation of two joints	
J.8	Polsterbind, ca. 10 cm x 2,7 m	
J.8	Padded bandages, approx. 10 cm x 2.7 m	
J.9	Gipsbind, ca. 12 cmx2m	
J.9	Plaster of paris bandage, approx. 12 cmx2m	
J.10	Fikseringsplaster, Tensoplast, ca. 2 cmx2m	
J.10	Adhesive tape, Tensoplast, approx. 2 cmx2m	
K.1	Båre, egnet til K.5	
K.1	Stretcher, suitable for K.5	
K.2	Oppustelig frakturskinne, senior, hel arm	
K.2	Inflatable fracture splint, senior, full arm	
K.3	Oppustelig frakturskinne, senior, hel ben	
K.3	Inflatable fracture splint, senior, full leg	
K.4	Stiv halskrave, indstillelig til forskellige positioner	
K.4	Stiff neck collar, adjustable	
K.5	Langt rygbræt, inkl. seler	
K.5	Spineboard, incl straps	
L.1	Coltosol®, færdigblandet tandcement i tuber à 8g	
L.1	Coltosol®, dental filling material ready for use, tubes of 8 g	
L.2	Skyllekanyle	
L.2	Dental rinse needle	
L.3	Specialinstrument til tandcement	
L.3	Special instrument for dental filling material	
L.4	Mundspejl	
L.4	Mouth mirror	
L.5	Ekskavator	
L.5	Excavator	
L.6	Fyldningsfjerner	
L.6	Filling remover	
L.7	Rustfri stålæske med låg til tandudlægeudstyr	
L.7	Instrument box, stainless steel with lid for dental equipment	

Grp. No	Medicinsk udstyr / Medical equipment	Mængde/ Quantity
M.1	Medicinmålebægre, ca. 25 ml, éngangs	
M.1	Graduated measuring cup, approx. 25 ml, disposable	
M.2	Varmedunk	
M.2	Hot-water bottle	
M.3	Bækken	
M.3	Bedpan	
N.1	Insektbekæmpelsesmiddel –pulver form	
N.1	Insecticide, powder form	
N.2	Kloreringsmiddel til drikkevand, 100 g	
N.2	Chlorinating agent for drinking water, 100 g	
N.3	Insektbekæmpelsesmiddel –væskeform	
N.3	Insecticide, liquid form	
O.2	Lægebog for Søfarende, udgivet af Seahealth (se note)	
O.2	Medical guide for Seafarers issued by Seahealth (see note)	
O.3	Fortegnelse, kontroldokument og brugervejledning kat. MFAG A2 (Download, www.soefartsstyrelsen.dk)	
O.3	Inventory, Control document and User instruction Cat. MFAG A2 (Download, www.dma.dk)	
O.5	Radio Medical Optegnelser	
O.5	Radio Medical Records	
O.6	MFAG (Medical First Aid Guide)	
O.6	MFAG (Medical First Aid Guide)	

O.2 Skal også forefindes på engelsk, såfremt der om bord er påmønstret ikke dansktalende besætningsmedlemmer.

O.2 Must also be available in English, if crewmembers who don't speak Danish have signed on.

O.3 skal være tilgængelig i digital eller printet udgave. www.soefartsstyrelsen.dk

O.3 must be available in digital or printed version. www.dma.dk

Liste ved aflevering af udløbet medicin til apotek
List when handing over expired medication to a pharmacy

Lægemiddelstof/ Active Pharmaceutical Ingredient	Adm. Form/ Form	Mængde / Amount
Morfin ^Δ	Inj. Væske	
Morphine ^Δ	Inj. Liqid	
Morfin ^Δ	Tablet / Kapsel	
Morphine ^Δ	Tablet / Capsula	
Cinnarizin ^Δ	Tablet	
Cinnarizine ^Δ	Tablets	
Diazepam ^Δ	Tablet	
Diazepam ^Δ	Tablets	
Diazepam ^Δ	Klyasma	
Diazepam ^Δ	Enema	
Haloperidol ^Δ	Inj. Væske	
Haloperidole ^Δ	Inj. Liquid	
Zolpidem ^Δ	Tablet	
Zolpidem ^Δ	Tablets	
Skib - kontrol nr. Ship - control no.	Skibsførerens Navn og Underskrift Masters Name and signature	
Dato / <i>Date</i>	Apotek / Pharmacy, eller/or Agent	
Navn / <i>Name</i>	Underskrift / <i>Signature</i>	
Stempel /stamp		

Ændringer / Changes

July-2024

Medicin som er fjernet / Medicine removed
3.5 Tbl. Tramadol 50 mg
7.4 Tbl. Erythromycin 500 mg
7.13 Tbl. Sulfamethoxazol+trimitroprim
8.3 Tbl. Quinin100 mg
10.4 Cream Sølvulfadiazin
12.5 Mixtur Laktulose 667 mg/ml
12.12 Tbl. Dopenridon10 mg
13.8 Dråber/gel Fusidin
15.2 Turbohaler Terbutalin
15.4 Turbohaler Budesonide
15.6 Inj. Salbutamol
16.6 Gelatine, Natriumchlorid

Medicin som er tilføjet / Medicine added
3.9 Tbl. Morfin 10 mg
4.6 Tbl. Prenisolon 25 mg
7.2 Tbl. Amoxicillin 500 mg
7.15 Tbl. Azithromycin 500mg
7.16 Tbl. Pivmecillinam 400 mg.
7.17 Tbl. Aciclovir 400 mg
8.4 Tbl. Artemether and lumefantrine
9.8 Tbl. Apixaban 5 mg
9.9 Inj. Tranexamsyre 100 mg/ml
10.8 Salve/ Ointment Fusidin 2%
12.13 Tbl Ondansetron 4 mg
12.14 Pulver/ powder Movicol®
13.13 Øjensalve / Ointment Chloramphenicol
15.7 Inhalation spray Sabumetol
15.8 Inhalation spray glycocortocoid
16.7 Steril vand / sterile water

Medicinsk udstyr som er fjernet / Medical equipment that is removed
E.4 Optiva*2 Intravenøs kanyle, 1,1 x 32, 20 g / <i>Optiva*2 intravenous cannula, 1.1 x 32, 20 g</i>
E.5 Vecafix® Fikseringsplaster til intravenøs kanyle / <i>Vecafix® Fixing plaster for intravenous cannula</i>

Medicinsk udstyr som er tilføjet / Medical equipment that is added
A.18 Hjertestarter (AED) 7/ <i>Automatic external defibrillator (AED)</i>
A.19 Iltbrille / <i>Nasal Cannula</i>
C.15 Absorberende underlag ca. 6 lag 60x40 cm, usteril, engangs / <i>Absorbent underlay, approx.6 layers, 60x40 cm, non-sterile, disposable</i>
C.16 Opkastpose / <i>Vomitbag</i>
D. 17 Ikke-absorberbar sutur, 5.0; 45 cm. FS eller DS25 / <i>Non-absorbable suture, 5.0; 45 cm. FS eller DS25</i>
E.4 Intravenøs kanyle, 1,1 x 32, 20 g (produkt navn fjernes) / <i>intravenous cannula, 1.1 x 32, 20 g (production name removed)</i>
E.5 Fikseringsplaster til intravenøs kanyle (produkt navn fjernes) / <i>Fixing plaster for intravenous cannula (product name removed)</i>
G.8 Hurtig test C-reaktiv protein (CRP) / <i>Rapid test for C- reactive proteins (CRP)</i>
I.1 Engangskatetre m. krum spids, CH 14, til mænd (antal øges til 10 stk.) / <i>Catheters for single use w/ curved tip, CH 14 for men (the amount increases to 10 pcs.)</i>

October 2023

4.1 Adrenalin opbevaring ændret - Læs indlægssedlen for opbevaring
Tilføjet liste aflevering af udløbet medicin på apotek

4.1 Epinephrine storage changed - read packet insert for storage
Added list of delivery of expired medicine at the pharmacy

July 2022

E.4 og E.5 Engelsk tekst rettet fra "needle" til cannula.
Generelle rettelser af stavfejl og layout

E.4 og E.5 Text corrected from "needle" to cannula.
General spelling corrections and layout

April 2022

B.14 rettet til B.18 Årepresse til massiv blødninger
Generelle rettelser af stavfejl og layout.

B.14 corrected to B.18 Tourniquet for massive bleeding
E.3 corrected to Tourniquet **for i.v.**
General spelling corrections and layout.

October 2021

Tilføjet Identifikation af skib (Side 1)
Tilføjet 13.12 Ciprofloxacin / Hydrocortison- øredråber
16.3 Hydroxyethylstivelse udgår
Tilføjet 16.6 Gelatine, Natrium chlorid erstatter 16.3
Tilføjet B.14 Årepresse til pulsåreblødning
Tilføjet C.13 Engangsforklæde
Tilføjet C.14 Sterilt lagen
Tilføjet F.7 Hypotermi termometer
Tilføjet I.3 Engangskatetre CH 14 til kvinder
Ændret N.1 Insektbekæmpelsesmiddel, ikke specificeret- **pulver form**
Tilføjet N.3 Insektbekæmpelsesmiddel, ikke specificeret- **væskeform**

Added Identification of ship (page 1)
Added 13.12 Ciprofloxacin / Hydrocortisone - eardrops
16.3 Hydroxyethyl amoid is discontinued product
Added 16.6 Gelatine sodium Chlorid replace 16.3
Added B.14 Tourniquet for massive bleeding
Added C.13 Disposable apron
Added C.14 Bed sheet, sterile
Added F.7 Hypotermi thermometer
Added I.3 Catheters for single use, CH 14 for women
Changed N.1 Insecticide, not specified - **powder form**
Added N.3 Insecticide, not specified - **liquid form**

July 2021

6.3 Diazepam klyasma styrke rettet tilbage til 4mg/ml - beklager ulejligheden
7.10 Vaccine Tetanus- label tilrettet
COVID-19 tillæg - overskrifter rettet. Tilføjet at hurtigtest skal være CE mærket

6.3 Diazepam enema strength corrected back to 4mg/ml - Sorry for the inconvenience
7.10 Vaccine Tetanus - label corrected
COVID-19 add on - the headlines corrected. Added that the quick test have to be CE marked.

February 2021

Tilføjet 8.4 Artemether og lumefantrine i "add medicine covid-19". Til behandling af Malaria.

Add. 8.4 Artemether and lumefantrine in "add medicine covid-19". For treatment of Malaria.

January 2021

8.2 Doxycyclin styrke ændret til 100 mg

8.2 Doxycyclin strength changed to 100 mg

Tilføjet side med ekstramedicin og medicinsk udstyr i forbindelse med Covid-19.

101-> Det er frivilligt om skibet vil medbringe medicinen og udstyr eller dele deraf.

Added page with additional medicine and medical equipment in connection with Covid-19. 101-> It is optional if the ship will bring the medicine and equipment or part of it.

Marts 2020

Ændringer er nu på sidste side

Changes is now a the last page

15.3 Inj. Terbutalin 0,5 mg/ml er erstattet med 15.6 Inj. Salbutamol 0,5 mg/ml

15.3 Inj. Terbutalin 0,5 mg/ml is changed to 15.6 Inj. Salbutamol 0,5 mg/ml

7.1 Phenoxymethylpenicillin styrken er ændret til 660 mg - 800 mg

7.1 Phenoxymethylpenicillin strenght change to 660 mg - 800 mg

D.4 Sutur fjernet firma navn

D.4 Sutur removed company name

August 2017

3.6 ændret til: 3.8 Diclofenac suppositorier 50 mg 50 stk.

7.10 ændret til tetanustoksid holdig vaccine 1. stk.

8.1 (Behandling) Brugervejledning rettet

O.2 Tilføjet ny lægebog for søfarende fra Seahealth

3.6 Changed to: 3.8 Diclofenac suppositorie 50 mg 50 pcs.

7.10 Changed to: Vaccine containing Tetanustoxid 1 pc.

8.1 (Treatment) User instruction corrected.

O.2 Add. the new Medical guide for seafarers from Seahealth